

Hardness Amplification and the Approximate Degree of Constant-Depth Circuits

Mark Bun ^{*}
mbun@seas.harvard.edu

Justin Thaler [†]
jthaler@fas.harvard.edu

Abstract

We establish a generic form of hardness amplification for the approximability of constant-depth Boolean circuits by polynomials. Specifically, we show that if a Boolean circuit cannot be pointwise approximated by low-degree polynomials to within constant error in a certain one-sided sense, then an OR of disjoint copies of that circuit cannot be pointwise approximated even with very high error. As our main application, we show that for every sequence of degrees $d(n)$, there is an explicit depth-three circuit $F : \{-1, 1\}^n \rightarrow \{-1, 1\}$ of polynomial-size such that any degree- d polynomial cannot pointwise approximate F to error better than $1 - \exp(-\tilde{\Omega}(nd^{-3/2}))$. As a consequence of our main result, we obtain an $\exp(-\tilde{\Omega}(n^{2/5}))$ upper bound on the discrepancy of a function in AC^0 , and an $\exp(\tilde{\Omega}(n^{2/5}))$ lower bound on the threshold weight of AC^0 , improving over the previous best results of $\exp(-\Omega(n^{1/3}))$ and $\exp(\Omega(n^{1/3}))$ respectively.

Our techniques also yield a new lower bound of $\Omega(n^{1/2}/\log^{(d-2)/2}(n))$ on the approximate degree of the AND-OR tree of depth d , which is tight up to polylogarithmic factors for any constant d , as well as new bounds for read-once DNF formulas. In turn, these results imply new lower bounds on the communication and circuit complexity of these classes, and demonstrate strong limitations on existing PAC learning algorithms.

1 Introduction

The ε -approximate degree of a Boolean function $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$, denoted $\widetilde{\deg}_\varepsilon(f)$, is the minimum degree of a real polynomial that approximates f to error ε in the ℓ_∞ norm. Approximate degree has pervasive applications in theoretical computer science. For example, lower bounds on approximate degree underly many tight lower bounds on quantum query complexity (e.g., [2, 4, 8, 28, 52]), and have been used to resolve several long-standing open questions in communication complexity [14, 16, 35, 50, 53, 54, 58, 59] (see the survey paper by Sherstov [47]). Meanwhile, upper bounds on approximate degree underly many of the fastest known learning algorithms, including PAC learning DNF and read-once formulas [6, 29], agnostically learning disjunctions [24], and PAC learning in the presence of irrelevant information [30, 44].

Despite the range and importance of these applications, large gaps remain in our understanding of approximate degree. The approximate degree of any *symmetric* Boolean function has been understood since Paturi's 1992 paper [41], but once we move beyond symmetric functions, few general results are known.

^{*}Harvard University, School of Engineering and Applied Sciences. Supported by an NDSEG Fellowship and NSF grant CNS-1237235.

[†]Simons Institute for the Theory of Computing at UC Berkeley. Parts of this work were performed while the author was a graduate student at Harvard University, School of Engineering and Applied Sciences. This work was supported by an NSF Graduate Research Fellowship, NSF grants CNS-1011840 and CCF-0915922, and a Research Fellowship from the Simons Institute for the Theory of Computing.

In this paper, we perform a careful study of the approximate degree of constant-depth Boolean circuits. In particular, we establish a generic form of hardness amplification for the pointwise approximation of small depth circuits by low-degree polynomials: we show that if a Boolean circuit f cannot be pointwise approximated to within constant error in a certain one-sided sense by polynomials of a given degree, then the circuit F obtained by taking an OR of disjoint copies of f cannot be approximated even with error exponentially close to 1. Notice that if f is computed by a circuit of polynomial size and constant depth, then so is F .

Our proof extends a recent line of work [15, 34, 45, 55] that seeks to prove approximate degree lower bounds by constructing explicit *dual polynomials*, which are dual solutions to a linear program that captures the approximate degree of any function. Specifically, we show that given a dual polynomial demonstrating that f cannot be approximated to within constant error, we can construct a dual polynomial demonstrating that F cannot be approximated even with error exponentially close to 1.

As the main application of our hardness amplification technique, for any $d > 0$ we exhibit an explicit function $F : \{-1, 1\}^n \rightarrow \{-1, 1\}$ computed by a polynomial size circuit of depth three for which any degree- d polynomial cannot pointwise approximate F to error $1 - \exp(-\tilde{\Omega}(nd^{-3/2}))$. We then use this result to obtain new bounds on two quantities that play central roles in learning theory, communication complexity, and circuit complexity: *discrepancy* and *threshold weight*. Specifically, we prove a new upper bound of $\exp(-\tilde{\Omega}(n^{2/5}))$ for the discrepancy of a function in AC^0 , and a new lower bound of $\exp(\tilde{\Omega}(n^{2/5}))$ for the threshold weight of AC^0 . As a second application, our hardness amplification result allows us to resolve, up to polylogarithmic factors, the approximate degree of AND-OR trees of arbitrary constant depth. Finally, our techniques also yield new lower bounds for read-once DNF formulas.

2 Overview of Results and Techniques

This section provides an overview of our results and the techniques we use to establish them. We defer detailed proofs to later sections.

2.1 Hardness Amplification

Recall that the ε -approximate degree of a Boolean function f is the minimum degree of a real polynomial that pointwise approximates f to error ε . Another fundamental measure of the complexity of f is its *threshold degree*, denoted $\text{deg}_{\pm}(f)$. The threshold degree of f is the least degree of a real polynomial that agrees in sign with f at all Boolean inputs.

Central to our results is a measure of the complexity of a Boolean function that we call *one-sided approximate degree*. This quantity, which we denote by $\widetilde{\text{odeg}}_{\varepsilon}(f)$, is an intermediate complexity measure that lies between ε -approximate degree and threshold degree. Unlike approximate degree and threshold degree, one-sided approximate degree treats inputs in $f^{-1}(1)$ and inputs in $f^{-1}(-1)$ asymmetrically.

More specifically, $\widetilde{\text{odeg}}_{\varepsilon}(f)$ captures the least degree of a *one-sided approximation* for f . Here, a one-sided approximation p for f is a polynomial that approximates f to error at most ε at all points $x \in f^{-1}(1)$, and satisfies the threshold condition $p(x) \leq -1 + \varepsilon$ at all points $x \in f^{-1}(-1)$. Notice that $\widetilde{\text{odeg}}_{\varepsilon}(f)$ is always *at most* $\widetilde{\text{deg}}_{\varepsilon}(f)$, but can be much smaller. Similarly, $\widetilde{\text{odeg}}_{\varepsilon}(f)$ is always *at least* $\text{deg}_{\pm}(f)$, but can be much larger.

One-sided approximate degree is the complexity measure that we amplify for constant-depth circuits: given a depth k circuit f on m variables that has one-sided approximate degree greater than d , we show how to generically transform f into a depth $k + 1$ circuit F on $t \cdot m$ variables such that F cannot be pointwise

approximated by degree d polynomials even to error $1 - 2^{-t}$.¹

Theorem 1. *Suppose $f : \{-1, 1\}^m \rightarrow \{-1, 1\}$ has one-sided approximate degree $\widetilde{\text{odeg}}_{1/2}(f) > d$. Denote by $F : \{-1, 1\}^{m \cdot t} \rightarrow \{-1, 1\}$ the block-wise composition $\text{OR}_t(f, \dots, f)$, where OR_t denotes the OR function on t variables. Then F cannot be pointwise approximated by degree- d polynomials even to within error $1 - 2^{-t}$ by degree- d polynomials. That is, the $(1 - 2^{-t})$ -approximate degree of F is greater than d .*

Remark: Theorem 1 demonstrates that $\widetilde{\text{odeg}}(f)$ admits a form of hardness amplification within AC^0 , which does not generally hold for the ordinary approximate degree. Indeed, Theorem 1 fails badly if the condition $\widetilde{\text{odeg}}_{1/2}(f) > d$ is replaced with the weaker condition $\widetilde{\text{deg}}_{1/2}(f) > d$ (in fact, $f = \text{OR}_m$ is a counterexample; see the discussion in Section 4 for details).

A dual formulation of one-sided approximate degree was previously exploited by Gavinsky and Sherstov to separate the multi-party communication versions of NP and co-NP [19], as well as by the current authors [15] and independently by Sherstov [45] to resolve the approximate degree of the two-level AND-OR tree. In this paper, we introduce the primal formulation of one-sided approximate degree, which allows us to express Theorem 1 as a hardness amplification result. We also argue for the importance of one-sided approximate degree as a Boolean function complexity measure in its own right.

Prior Work on Hardness Amplification for Approximate Degree. For the purposes of this discussion, we informally consider a hardness amplification result for approximate degree to be any statement of the following form: Fix two functions $f : \{-1, 1\}^m \rightarrow \{-1, 1\}$ and $g : \{-1, 1\}^t \rightarrow \{-1, 1\}$. Then the composed function $g(f, \dots, f) : \{-1, 1\}^{m \cdot t} \rightarrow \{-1, 1\}$ is strictly harder to approximate in the ℓ_∞ norm by low-degree polynomials than is the original function f .

We think of such a result as establishing that application of the outer function g to t disjoint copies of f amplifies the hardness of f . Here we consider polynomial degree to be a resource, and “harder to approximate” can refer either to the amount of resources required for the approximation, to the error of the approximation, or to a combination of the two.

Two particular kinds of hardness amplification results for approximate degree have received particular attention. *Direct-sum* theorems focus on amplifying the degree required to obtain an approximation, but do not focus on amplifying the error. For example, a typical direct-sum theorem identifies conditions on f and g that guarantee that $\widetilde{\text{deg}}_\epsilon(g(f, \dots, f)) \geq \widetilde{\text{deg}}_\epsilon(g) \cdot \widetilde{\text{deg}}_\epsilon(f)$. In contrast, a *direct-product* theorem focuses on amplifying both the error and the minimum degree required to achieve this error. An *XOR lemma* is a special case of either type of theorem where the combining function g is the XOR function. Ideally, an XOR lemma of the direct-product form establishes that there exists a sufficiently small constant $\delta > 0$ such that $\widetilde{\text{deg}}_{1-2^{-\delta t}}(\text{XOR}_t(f, \dots, f)) \geq t \cdot \widetilde{\text{deg}}_{1/3}(f)$. That is, an XOR lemma establishes that approximating the XOR of t disjoint copies of f requires a t -fold blowup in degree relative to f , even if one allows error exponentially close to 1.

O’Donnell and Servedio [40] proved an XOR lemma for *threshold degree*, establishing that $\text{XOR}_t(f, \dots, f)$ has threshold degree t times the threshold degree of f . In later work, Sherstov [55] proved a direct sum result for approximate degree that holds whenever the combining function g has low block-sensitivity. His techniques also capture O’Donnell and Servedio’s XOR lemma for threshold degree as a special case. In [52], Sherstov proved a number of hardness amplification results for approximate degree. Most notably, he proved an optimal XOR lemma, as well as a direct-sum theorem that holds whenever the combining function has close to maximal approximate degree (i.e., approximate degree $\Omega(t)$). Sherstov used his XOR lemma to prove direct product theorems for quantum query complexity, and in subsequent work [53], to show direct product theorems for the multiparty communication of set disjointness.

¹Follow-up work by Sherstov [46] has established a lower bound on the *threshold degree* of F . Specifically, he has shown that there is some constant c such that $\text{deg}_\pm(F) > \min\{ct, d\}$. See Section 2.5 for further discussion of this result.

Comparison to Prior Work. In this paper, we are interested in establishing approximate degree lower bounds for constant-depth circuits over the basis $\{\text{AND}, \text{OR}, \text{NOT}\}$. For this purpose, it is essential to consider combining functions (such as OR, see Theorem 1) that are themselves in AC^0 , ruling out the use of XOR as a combining function. Our hardness amplification result (Theorem 1) is orthogonal to direct-sum theorems: direct-sum theorems focus on amplifying degree but not error, while Theorem 1 focuses on amplifying error but not degree. Curiously, Theorem 1 is nonetheless a critical ingredient in our proof of a direct-sum type theorem for AND-OR trees of constant depth (Theorem 8).

Proof Idea. As discussed in the introduction, our proof of Theorem 1 relies on a dual characterization of one-sided approximate degree (cf. Theorem 15). Specifically, for any m -variate Boolean function f satisfying $\widetilde{\text{odeg}}_{1/2}(f) > d$, there exists a dual object $\psi : \{-1, 1\}^m \rightarrow \mathbb{R}$ that witnesses this fact — we refer to ψ as a “dual polynomial” for f . As we show in Theorem 15, ψ satisfies three important properties: (1) ψ has high correlation with f , (2) ψ has zero correlation with all polynomials of degree at most d , and (3) $\psi(x)$ agrees in sign with $f(x)$ for all $x \in f^{-1}(-1)$. We refer to the second property by saying that ψ has *pure high degree* d , and we refer to the third property by saying that ψ has *one-sided error*.

Our proof proceeds by taking a dual witness ψ to the high one-sided approximate degree of f , and a certain dual witness Ψ for the function OR_t , and combining them to obtain a dual witness ζ for the fact that $\widetilde{\text{deg}}_{1-2^{-t}}(\text{OR}_t(f, \dots, f)) > d$. Our analysis of the combined dual witness crucially exploits two properties: first, that ψ has one-sided error and second, that the vector whose entries are all equal to -1 has very large (in fact, maximal) Hamming distance from the unique input in $\text{OR}_t^{-1}(1)$.

Our method of combining the two dual witnesses was first introduced by Sherstov [55, Theorem 3.3] and independently by Lee [34]. This method has also been exploited by the present authors in [15] to resolve the approximate degree of the two-level AND-OR tree, and by Sherstov [52] to prove direct sum and direct product theorems for polynomial approximation. However, as discussed above, prior work used this method of combining dual witnesses exclusively to amplify the *degree* in the resulting lower bound; in contrast, we use the combining method in the proof of Theorem 1 to amplify the *error* in the resulting lower bound.

From a technical perspective, the primary novelty in the proof of Theorem 1 lies in our choice of an appropriate (and simple) dual witness Ψ for OR_t , and the subsequent analysis of the correlation of the combined witness ζ with $\text{OR}_t(f, \dots, f)$. By our choice of Ψ , we are able to show that ζ has correlation with $\text{OR}_t(f, \dots, f)$ that is *exponentially* close to 1, yielding a lower bound even on the degree of approximations with very high error.

2.2 Lower Bounds For AC^0

2.2.1 A New One-Sided Approximate Degree Lower Bound for AC^0

Our ultimate goal is to use Theorem 1 to construct a function F in AC^0 that is hard to approximate by low-degree polynomials even with error exponentially close to 1. However, in order to apply Theorem 1, we must first identify an AC^0 function f such that $\widetilde{\text{odeg}}_{1/2}(f)$ is large.

To this end, we identify fairly general conditions guaranteeing that the one-sided approximate degree of a function is *equal* to its approximate degree, up to a logarithmic factor. To express our result, let $[N] = \{1, \dots, N\}$, and let m, N, R be a triple of positive integers such that $R \geq N$, and $m = N \cdot \log_2 R$. In most cases, we will take $R = N$. We specifically consider Boolean functions f on $\{-1, 1\}^m$ that interpret their input x as the values of a function g_x mapping $[N] \rightarrow [R]$. That is, we break x up into N blocks each of length $\log_2 R$, and regard each block x_i as the binary representation of $g_x(i)$. Hence, we think of f as computing some *property* ϕ_f of functions $g_x : [N] \rightarrow [R]$. We say that a property ϕ is *symmetric* if for all $g : [N] \rightarrow [R]$, all permutations σ on $[R]$, and all permutations π on $[N]$, it holds that $\phi(g) = \phi(\sigma \circ g \circ \pi)$.

Theorem 2. *Let $f : \{-1, 1\}^m \rightarrow \{-1, 1\}$ be a Boolean function corresponding to a symmetric property ϕ_f of functions $g_x : [N] \rightarrow [R]$. Suppose that for every pair $x, y \in f^{-1}(-1)$, there is a pair of permutations σ on $[R]$ and π on $[N]$ such that $g_x = \sigma \circ g_y \circ \pi$. Then $\widetilde{\text{odeg}}_\varepsilon(f) \geq \frac{1}{\log_2 R} \cdot \widetilde{\text{deg}}_\varepsilon(f)$ for all $\varepsilon > 0$.*

Proof Idea. It is enough to show that any one-sided ε -approximation p to f can be transformed into an actual ε -approximation r to f in a manner that does not increase the degree by too much (i.e., in a manner guaranteeing that $\text{deg}(r) \leq (\log_2 R) \text{deg}(p)$).

Our transformation from p to r consists of two steps. In the first step, we turn p into a “symmetric” polynomial $p^{\text{sym}}(x) := \mathbb{E}_{y \sim x}[p(y)]$ where $y \sim x$ if $g_y = \sigma \circ g_x \circ \pi$ for some permutations σ on $[R]$ and π on $[N]$. It follows from work of Ambainis [4] (see Lemma 23) that the map $p \mapsto p^{\text{sym}}$ increases the degree of p by a factor of at most $\log_2 R$. In the second step, we argue that there is an affine transformation r of p^{sym} that is an actual ε -approximation to f , completing the construction.

The existence of the affine transformation r of p^{sym} follows from two observations: (1) if p is a one-sided approximation for f , then so is p^{sym} (this holds because ϕ_f is symmetric), and (2) p^{sym} takes on a constant value v on $f^{-1}(-1)$, i.e., $p^{\text{sym}}(x) = v$ for all $x \in f^{-1}(-1)$ (this holds because $x \sim y$ for every pair of inputs $x, y \in f^{-1}(-1)$). These observations imply that even if p^{sym} is a very poor approximation to f on $f^{-1}(-1)$, we can still obtain a good pointwise approximation r by applying an affine transformation to the range of p^{sym} that maps v to -1 and moves all values closer to 1. Section 5.1 contains the details.

In our primary application of Theorem 2, we let $f : \{-1, 1\}^m \rightarrow \{-1, 1\}$ be the ELEMENT DISTINCTNESS function (defined in Section 3). Aaronson and Shi [2] showed that the approximate degree of ELEMENT DISTINCTNESS is $\Omega((m/\log m)^{2/3})$. ELEMENT DISTINCTNESS is computed by a CNF of polynomial size, and Aaronson and Shi’s result remains essentially the best-known lower bound for the approximate degree of a function in AC^0 . Theorem 2 applies to ELEMENT DISTINCTNESS, yielding the following corollary.

Corollary 3. *Let $f : \{-1, 1\}^m \rightarrow \{-1, 1\}$ denote the ELEMENT DISTINCTNESS function. Then $\widetilde{\text{odeg}}(f) = \tilde{\Omega}(m^{2/3})$.*

The best known lower bound on the one-sided approximate degree of an AC^0 function that followed from prior work was $\Omega(m^{1/2})$ (which holds for the AND function [19, 39], cf. Fact 16 in Section 3). Section 2.5 describes some further implications of Theorem 2.

Remark: In an earlier version of this work, we gave a different *dual* proof of Corollary 3. Specifically we showed (cf. Appendix C) that any dual witness for the high approximate degree of ELEMENT DISTINCTNESS can be transformed into a dual witness with one-sided error. This proof in fact shows that $\widetilde{\text{odeg}}(f) = \widetilde{\text{deg}}(f)$ (i.e. without incurring the loss of a $1/\log_2 R$ factor as in Theorem 2). However it remains unclear how to generalize this dual argument to the more general class of properties to which Theorem 2 applies (including the 2-TO-1 property discussed in Section 2.5 below). Theorem 2 therefore provides an example of a setting in which the primal view of one-sided approximate degree introduced in this work may be easier to reason about than the dual formulation used in prior work.

2.2.2 Accuracy-Degree Tradeoff Lower Bounds for AC^0

By Corollary 3, we can apply Theorem 1 to ELEMENT DISTINCTNESS to obtain a depth-three Boolean circuit F with $t \cdot m$ inputs such that $\widetilde{\text{deg}}_\varepsilon(F) = \tilde{\Omega}(m^{2/3})$, for $\varepsilon = 1 - 2^{-t}$. By choosing t and m appropriately, we obtain a depth-three circuit on $n = t \cdot m$ variables of size $\text{poly}(n)$ such that any degree- d polynomial cannot pointwise approximate F to error better than $1 - \exp\left(-\tilde{\Omega}(nd^{-3/2})\right)$.

Reference	Discrepancy Bound	Circuit Depth
Sherstov [51]	$\exp(-\Omega(n^{1/5}))$	3
Buhrman et al. [14]	$\exp(-\Omega(n^{1/3}))$	3
Sherstov [50]	$\exp(-\Omega(n^{1/3}))$	3
This work	$\exp(-\tilde{\Omega}(n^{2/5}))$	4

Table 1: Comparison of our new discrepancy bound for AC^0 to prior work. The circuit depth column lists the depth of the circuit used to exhibit the bound.

Corollary 4. *For every $d > 0$, there is a depth-3 Boolean circuit $F : \{-1, 1\}^n \rightarrow \{-1, 1\}$ of size $\text{poly}(n)$ such that any degree- d polynomial cannot pointwise approximate F to error better than $1 - \exp(-\tilde{\Omega}(nd^{-3/2}))$. In particular, there is a depth-3 circuit F such that any polynomial of degree at most $n^{2/5}$ cannot pointwise approximate F to error better than $1 - \exp(-\tilde{\Omega}(n^{2/5}))$.*

2.2.3 Discrepancy Upper Bound

Discrepancy, defined formally in Section 5.4, is a central quantity in communication complexity and circuit complexity. For instance, upper bounds on the discrepancy of a function f immediately yield lower bounds on the cost of small-bias communication protocols for computing f (Section 8 has details). The first exponentially small discrepancy upper bounds for AC^0 were proved by Burhman et al. [14] and Sherstov [50, 51], who exhibited constant-depth circuits with discrepancy $\exp(-\Omega(n^{1/3}))$. Our results improve the best-known upper bound to $\exp(-\tilde{\Omega}(n^{2/5}))$.

Our result relies on a powerful technique developed by Sherstov [50], known as the pattern-matrix method (stated as Theorem 26 in Section 5.4). This technique allows one to automatically translate lower bounds on the ε -approximate degree of a Boolean function F into lower bounds on the *discrepancy* of a related function F' as long as ε is exponentially close to one. By applying the pattern-matrix method to Corollary 4, we obtain the following result.

Corollary 5. *There is a depth-4 Boolean circuit $F' : \{-1, 1\}^n \rightarrow \{-1, 1\}$ with discrepancy $\exp(-\tilde{\Omega}(n^{2/5}))$.*

2.2.4 Threshold Weight Lower Bound

A *polynomial threshold function* (PTF) for a Boolean function f is a multilinear polynomial p with integer coefficients that agrees in sign with f on all Boolean inputs. The *weight* of an n -variate polynomial p is the sum of the absolute value of its coefficients. The *degree- d threshold weight* of a Boolean function $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$, denoted $W(f, d)$, refers to the least weight of a degree- d PTF for f . We let $W(f)$ denote the quantity $W(f, n)$, i.e., the least weight of any threshold function for f regardless of its degree. As discussed in Section 8, threshold weight has important applications in learning theory.

Threshold weight is closely related to ε -approximate degree when ε is very close to 1 (see Lemma 20 in Section 3.2). This allows us to translate Corollary 4 into a lower bound on the degree- d threshold weight of AC^0 .

Corollary 6. *For every $d > 0$, there is a depth-3 Boolean circuit $F : \{-1, 1\}^n \rightarrow \{-1, 1\}$ of size $\text{poly}(n)$ such that $W(F, d) \geq \exp(\tilde{\Omega}(nd^{-3/2}))$. In particular, $W(F, n^{2/5}) = \exp(\tilde{\Omega}(n^{2/5}))$.*

A result of Krause [32] (see Lemma 27 in Section 5.5) allows us to extend our new degree- d threshold weight lower bound for F into a *degree independent* threshold weight lower bound for a related function

F' (we also provide a new and simple proof of Krause’s result based on LP duality, cf. Appendix D). The previous best lower bound on the threshold weight of AC^0 was $\exp(\Omega(n^{1/3}))$, due to Krause and Pudlák [33].

Corollary 7. *There is a depth-4 Boolean circuit $F' : \{-1, 1\}^n \rightarrow \{-1, 1\}$ satisfying $W(F') = \exp(\tilde{\Omega}(n^{2/5}))$.*

Moreover, while the threshold weight bound of Corollary 7 is stated for polynomial threshold functions over $\{-1, 1\}^n$, we show that the same threshold weight lower bound also holds for polynomials over $\{0, 1\}^n$.

2.3 Approximate Degree Lower Bounds for AND-OR Trees

The d -level AND-OR tree on n variables is a function described by a read-once circuit of depth d consisting of alternating layers of AND gates and OR gates. We assume for simplicity that all gates have fan-in $n^{1/d}$. For example, the two-level AND-OR tree is a read-once CNF in which all gates have fan-in $n^{1/2}$.

Until recently, the approximate degree of AND-OR trees of depth two or greater had resisted characterization, despite 19 years of attention [4, 15, 22, 39, 45, 55, 57]. The case of depth two was reposed as a challenge problem by Aaronson in 2008 [1], as it captured the limitations of existing lower bound techniques. This case was resolved last year by the current authors [15], and independently by Sherstov [45], who proved a lower bound of $\Omega(\sqrt{n})$, matching an upper bound of Høyer, Mosca, and de Wolf [22]. However, the case of depth three or greater remained open. To our knowledge, the best known lower bound for $d \geq 3$ was $\Omega(n^{1/4+1/2d})$, which follows by combining the depth-two lower bound [15, 45] with an earlier direct-sum theorem of Sherstov [55, Theorem 3.1].

By combining the techniques of our earlier work [15] with our hardness amplification result (Theorem 1), we improve this lower bound to $\Omega(n^{1/2}/\log^{(d-2)/2}(n))$ for any constant $d \geq 2$. A line of work on quantum query algorithms [6, 22, 43] established an upper bound of $O(n^{1/2})$ for AND-OR trees of any depth, demonstrating that our result is optimal up to polylogarithmic factors (see Section 6 for details).

Theorem 8. *Let $\text{AND-OR}_{d,n}$ denote the d -level AND-OR tree on n variables. Then $\widetilde{\text{deg}}(\text{AND-OR}_{d,n}) = \Omega(n^{1/2}/\log^{(d-2)/2} n)$ for any constant $d > 0$.*

Proof Idea. To introduce our proof technique, we first describe the method used in [15] to construct an optimal dual polynomial in the case $d = 2$, and we identify why this method breaks down when trying to extend to the case $d = 3$. We then explain how to use our hardness amplification result (Theorem 1) to construct a different dual polynomial that does extend to the case $d = 3$.

Let M denote the fan-in of all gates in $\text{OR-AND}_{2,M^2}$. In our earlier work [15], we constructed a dual polynomial for $\text{OR-AND}_{2,M^2}$ as follows.² By Fact 16 there is a dual polynomial γ_1 witnessing the fact that $\widetilde{\text{odeg}}(\text{AND}_M) = \Omega(M^{1/2})$, and a dual polynomial γ_2 witnessing the fact that $\widetilde{\text{deg}}(\text{OR}_M) = \Omega(M^{1/2})$. We then combined the dual witnesses γ_1 and γ_2 , using the same “combining” technique as in the proof of Theorem 1, to obtain a dual witness $\gamma_3 : \{-1, 1\}^{M^2} \rightarrow \mathbb{R}$ for the high approximate degree of $\text{OR-AND}_{2,M^2}$.

Recall that we say a dual witness has *pure high degree* d if it has zero correlation with every polynomial of degree at most d . It followed from earlier work [55] that γ_3 has pure high degree equal to the product of the pure high degree of γ_1 and the pure high degree of γ_2 , yielding an $\Omega(M)$ lower bound on the pure high degree of γ_3 . The new ingredient of the analysis in [15] was to use the one-sided error of the “inner” dual witness γ_1 to argue that γ_3 also had good correlation with $\text{OR-AND}_{2,M^2}$.

²We actually constructed a dual polynomial for $\text{AND-OR}_{2,M^2}$, but the analysis for the case of $\text{OR-AND}_{2,M^2}$ is entirely analogous.

Extending to Depth Three. Let $M = n^{1/3}$ denote the fan-in of all gates in $\text{AND-OR}_{3,n}$. In constructing a dual witness for $\text{AND-OR}_{3,n} = \text{AND}_M(\text{OR-AND}_{2,M^2}, \dots, \text{OR-AND}_{2,M^2})$, it is natural to try the following approach. Let γ_4 be a dual polynomial witnessing the fact that the approximate degree of $\text{AND}_M = \Omega(\sqrt{M})$. Then we can combine γ_3 and γ_4 in the same manner as above to obtain a dual function γ_5 .

The difficulty in establishing that γ_5 is a dual witness to the high approximate degree of $\text{AND-OR}_{3,n}$ is in showing that γ_5 has good correlation with AND-OR_3 . In our earlier work, we showed γ_3 has large correlation with $\text{OR-AND}_{2,n}$ by exploiting the fact that the inner dual witness γ_1 had one-sided error, i.e., $\gamma_1(y)$ agrees in sign with AND_M whenever $y \in \text{AND}_M^{-1}(-1)$. However, γ_3 itself does not satisfy an analogous property: there are inputs $x_i \in \text{OR-AND}_{2,M^2}^{-1}(-1)$ such that $\gamma_3(x_i) > 0$, and there are inputs $x_i \in \text{OR-AND}_{2,M^2}^{-1}(1)$ such that $\gamma_3(x_i) < 0$.

To circumvent this issue, we use a different inner dual witness γ'_3 in place of γ_3 . Our construction of γ'_3 utilizes our hardness amplification analysis to achieve the following: while γ'_3 has error “on both sides”, the error from the “wrong side” is very small. The hardness amplification step causes γ'_3 to have pure high degree that is lower than that of the dual witness γ_3 constructed in [15] by a $\sqrt{\log n}$ factor. However, the hardness amplification step permits us to prove the desired lower bound on the correlation of γ_5 with $\text{AND-OR}_{3,n}$. The proof for the general case, which is quite technical, can be found in Section 6.

2.4 Lower Bounds for Read-Once DNFs and CNFs

Our techniques also yield new lower bounds on the approximate degree and degree- d threshold weight of read-once DNF and CNF formulas. Before stating our results, we discuss relevant prior work.

In their seminal work on perceptrons, Minsky and Papert exhibited a read-once DNF $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$ with *threshold degree* $\Omega(n^{1/3})$ [37]. That is, a real polynomial requires degree $\Omega(n^{1/3})$ just to agree with f in sign. However, to our knowledge no non-trivial lower bound on the degree- d threshold *weight* of read-once DNFs was known for any $d = \omega(n^{1/3})$.

In an influential result, Beigel [11] exhibited a polynomial-size (read-many) DNF called ODD-MAX-BIT satisfying the following: there is some constant $\delta > 0$ such that $\deg_{1-2^{-\delta n/d^2}}(\text{ODD-MAX-BIT}) > d$, and hence also $W(\text{ODD-MAX-BIT}, d) = \exp(\Omega(n/d^2))$ (see Section 3.2). Motivated by applications in computational learning theory (see Section 8), Klivans and Servedio showed that Beigel’s lower bound is essentially tight for $d < n^{1/3}$ [30]. Very recently, Servedio, Tan, and Thaler showed an alternative lower bound on the degree- d threshold weight of ODD-MAX-BIT. Specifically, they showed that $W(\text{ODD-MAX-BIT}, d) = \exp\left(\Omega\left(\sqrt{n/d}\right)\right)$ [44]. The lower bound of Servedio et al. improves over Beigel’s for any $d > n^{1/3}$, and is essentially tight in this regime (i.e., when $d > n^{1/3}$).

While ODD-MAX-BIT is a relatively simple DNF (in fact, it is a *decision list*), it is not a read-once DNF. Our results extend the lower bounds of Servedio et al. and Beigel from decision lists to read-once DNFs and CNFs. In the statement of the results below, we restrict ourselves to DNFs, as the case of CNFs is entirely analogous.

2.4.1 Extending the Lower Bound of Servedio et al. to Read-Once DNFs

In order to extend the lower bound of Servedio et al. to read-once DNFs and CNFs, we extend our hardness amplification techniques from one-sided approximate degree to a new quantity we call *degree- d one-sided non-constant approximate weight*. This quantity captures the least L_1 *weight* (excluding the constant term) of a polynomial of degree at most d that is a one-sided approximation of f . We denote the degree- d one-sided approximate weight of a Boolean function f by $W_\varepsilon^*(f, d)$, where ε is an error parameter.

We prove the following analog of Theorem 1.

Theorem 9. Fix $d > 0$. Let $f : \{-1, 1\}^m \rightarrow \{-1, 1\}$, and suppose that $W_{3/4}^*(f, d) > w$. Let $F : \{-1, 1\}^{m \cdot t} \rightarrow \{-1, 1\}$ denote the function $\text{OR}_t(f, \dots, f)$. Then any degree- d polynomial that approximates F to within error $1 - 2^{-t}$ requires weight $2^{-5t}w$.

Adapting a proof of Servedio et al., we can show that $W_{3/4}^*(\text{AND}_m, d) \geq 2^{\Omega(m/d)}$. By applying Theorem 9 with $f = \text{AND}_m$, along with standard manipulations, we are able to extend the lower bound of Servedio et al. to read-once CNFs and DNFs.

Corollary 10. For each $d = o(n/\log^4 n)$, there is a read-once DNF F satisfying $W(F, d) = \exp\left(\Omega\left(\sqrt{n/d}\right)\right)$.

In particular, there is a read-once DNF that cannot be computed by any PTF of $\text{poly}(n)$ weight, unless the degree is $\tilde{\Omega}(n)$.

2.4.2 Extending Beigel’s Lower Bound to Read-Once DNFs

It is known that $\widetilde{\text{odeg}}(\text{AND}_m) = \Omega(m^{1/2})$ (cf. Fact 16). By applying Theorem 1 with $f = \text{AND}_m$, we obtain the following result.

Corollary 11. There is an (explicit) read-once DNF $F : \{-1, 1\}^n \rightarrow \{-1, 1\}$ with $\widetilde{\text{deg}}_{1-2^{-n/d^2}}(F) = \Omega(d)$.

We remark that for $d < n^{1/3}$, Corollary 11 is subsumed by Minsky and Papert’s seminal result that exhibited a read-once DNF F with threshold degree $\Omega(n^{1/3})$ [37]. However, for $d > n^{1/3}$, it is not subsumed by Minsky and Papert’s result, nor by Corollary 10. Indeed, Corollary 10 yields a lower bound on the degree- d threshold weight of read-once DNFs, but not a lower bound on the *approximate-degree* of read-once DNFs (see Section 3.2 for further discussion on the separation between these quantities).

2.5 Discussion

2.5.1 Subsequent Work by Sherstov

In 1969, Minsky and Papert gave a lower bound of $\Omega(n^{1/3})$ on the threshold degree of an explicit read-once DNF formula. Klivans and Servedio [29] proved their lower bound to be tight within a logarithmic factor for DNFs of polynomial size, but it remained a well-known open question to give a threshold degree lower bound of $\Omega(n^{1/3+\delta})$ for a function in AC^0 ; the only progress prior to our work was due to O’Donnell and Servedio [40], who established an $\Omega(n^{1/3} \log^k n)$ lower bound for any constant $k > 0$.

Let f denote the ELEMENT DISTINCTNESS function on $n^{3/5}$ variables. In an earlier version of this work, we conjectured that the function $F = \text{OR}_{n^{2/5}}(f, \dots, f)$ appearing in Corollary 4 in fact satisfies $\text{deg}_{\pm}(f) = \tilde{\Omega}(n^{2/5})$, and observed that this would yield the first polynomial improvement on Minsky and Papert’s lower bound. Sherstov [46, Theorem 7.1] has recently proved our conjecture. His proof, short and elegant, extends our dual witness construction in the proof of Theorem 1 to establish a different form of hardness amplification, from one-sided approximate degree to threshold degree. Specifically, he shows that if a Boolean function f has one-sided approximate degree d , then the block-wise composition $\text{OR}_t(f, \dots, f)$ has threshold degree at least $\min\{ct, d\}$ for some constant c . This result is incomparable to our Theorem 1 when $t \leq d$, but when $t \gg d$, Sherstov’s result is a substantial strengthening of Theorem 1.

In the same work, Sherstov has also proven a much stronger and more difficult result: for any $k > 2$, he gives a read-once formula of depth k with threshold degree $\Omega(n^{(k-1)/(2k-1)})$. Notice that for any constant $\delta > 0$, this yields an AC^0 function with threshold degree $\Omega(n^{1/2-\delta})$. This in turn yields an improvement of our discrepancy upper bound (Corollary 5) for AC^0 to $\exp(-\Omega(n^{1/2-\delta}))$, and of our threshold weight lower bound (Corollary 7) to $\exp(\Omega(n^{1/2-\delta}))$.

2.5.2 Subsequent Work by Kanade and Thaler

Existing applications of one-sided approximate degree [15, 19, 45, 46] have all been of a negative nature (proving communication and circuit lower bounds, establishing limitations on existing PAC learning algorithms, etc.). Kanade and Thaler [25] have recently identified a positive (algorithmic) application of one-sided approximate degree. Specifically, they show that one-sided approximate degree upper bounds imply fast algorithms in the reliable agnostic learning framework of Kalai et al. [23]. This framework captures learning tasks in which one type of error (such as false negative errors) is costlier than other types. Kanade and Thaler use this result to give the first sub-exponential time algorithms for distribution-independent reliable learning of several fundamental concept classes.

In light of these developments, we are optimistic that the notion of one-sided approximate degree will continue to enable progress on questions within the analysis of Boolean functions and computational complexity theory.

2.5.3 Future Directions

Beame and Machmouchi [9] established an $\Omega(n/\log n)$ lower bound on the quantum query complexity of a specific function $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$ in AC^0 . The previous best lower bound was $\Omega((n/\log n)^{2/3})$, which held for the ELEMENT DISTINCTNESS function [2].

Beame and Machmouchi's lower bound applies to the 2-TO-1 function, which is computed by depth-three circuit of polynomial size. This function interprets its input as a list of N numbers from a range of size $R \geq N$, and evaluates to -1 if and only if exactly $N/2$ numbers appear in the list, each with frequency exactly 2. They pose as an open question the problem of resolving the approximate degree of 2-TO-1³ (recall that the approximate degree of f is a lower bound on the quantum query complexity of f , but polynomial separations between approximate degree and quantum query complexity are known [5]).

For simplicity, we focus on the case where $R = N$. We observe that Theorem 2 applies to the 2-TO-1 function, revealing that its one-sided approximate degree is almost equal to its approximate degree.

Corollary 12. *Let $f : \{-1, 1\}^m \rightarrow \{-1, 1\}$ denote the 2-TO-1 function on m variables. For any $\varepsilon > 0$, $\widetilde{\text{odeg}}_\varepsilon(f) \geq \widetilde{\text{deg}}_\varepsilon(f)/\log m$.*

Combining Corollary 12 and the recent result [46, Theorem 7.1] allows us to transform any *approximate degree* lower bound for the 2-TO-1 function into a *threshold degree* lower bound for a related depth-four circuit.

Corollary 13. *Let $f : \{-1, 1\}^m \rightarrow \{-1, 1\}$ denote the 2-TO-1 function on m variables, and let $d = \widetilde{\text{odeg}}(f) \geq \widetilde{\text{deg}}(f)/\log m$. Let $n = m \cdot d$, and define $F : \{-1, 1\}^n \rightarrow \{-1, 1\}$ via $F = \text{OR}_d(f, \dots, f)$. Then $\text{deg}_\pm(F) = \Omega(d)$. In particular, if $\widetilde{\text{deg}}(f) = \Omega(m/\log m)$, then $\text{deg}_\pm(F) = \Omega(n^{1/2}/\log n)$.*

Thus, establishing a quasilinear lower bound on the approximate degree of 2-TO-1 would immediately yield a function F computable by a depth-four circuit of polynomial size with threshold degree $\tilde{\Omega}(n^{1/2})$, a polynomial improvement over Sherstov's $\Omega(n^{(k-1)/(2k-1)})$ bound for any constant depth k . Even a lower bound of $\Omega(m^{3/4+\delta})$ for some positive constant δ on the approximate degree of the 2-TO-1 function would yield a depth four circuit with threshold degree $\Omega(n^{3/7+\delta'})$ for some $\delta' > 0$. This would constitute a polynomial improvement over the current best lower bound of $\Omega(n^{3/7})$ for depth 4, and would additionally imply improved lower bounds on the threshold weight and discrepancy of depth five circuits.

³Technically speaking, they ask about the 2-TO-1-VS-ALMOST-2-TO-1 function, which is a promise variant of the 2-TO-1 function.

While the best known lower bound on the approximate degree of the 2-TO-1 function on m variables is currently $\Omega((m/\log m)^{2/3})$ (this can be derived by reduction to ELEMENT DISTINCTNESS), we conjecture that its approximate degree is in fact $\Omega(m/\log m)$, and interpret Beame and Machmouchi’s quantum query lower bound as providing mild evidence in favor of this conjecture.

2.6 Paper Roadmap

Section 3 establishes terminology, introduces our main technique based on LP-duality, and proves essential technical lemmas. Section 4 establishes our central hardness amplification result for one-sided approximate degree (Theorem 1). Section 5 establishes our new one-sided approximate degree lower bound for AC^0 (Theorem 2, Corollary 3), and combines this with Theorem 1 to obtain our new lower bounds on “accuracy vs. degree” tradeoffs for pointwise approximating AC^0 by polynomials (Corollary 4). It then proves our new discrepancy upper bound for an AC^0 function (Corollary 5) and our new threshold weight lower bound for AC^0 (Corollaries 6 and 7). Section 6 proves our new approximate degree lower bound for AND-OR trees (Theorem 8). Section 7 proves our new lower bounds for read-once DNFs (Theorem 9, Corollary 10, and Corollary 11). Section 8 highlights several applications of these results to communication complexity, circuit lower bounds, and learning theory.

3 Preliminaries

We work with Boolean functions $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$ under the standard convention that 1 corresponds to logical false, and -1 corresponds to logical true. For a real-valued function $r : \{-1, 1\}^n \rightarrow \mathbb{R}$, we let $\|r\|_\infty = \max_{x \in \{-1, 1\}^n} |r(x)|$ denote the ℓ_∞ norm of r . We let OR_n and AND_n denote the OR function and AND function on n variables respectively. Define $\widetilde{\text{sgn}}(t) = -1$ if $t \leq 0$ and 1 otherwise. For a set $S \subseteq [n] = \{1, \dots, n\}$, let $\chi_S(x) := \prod_{i \in S} x_i$ denote the parity function over variables indexed by S .

We now define the notions of approximate degree, approximate weight, threshold degree, threshold weight, and their one-sided variants.

3.1 Polynomial Approximations and their Dual Characterizations

3.1.1 Approximate Degree

The ε -approximate degree of a function $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$, denoted $\widetilde{\text{deg}}_\varepsilon(f)$, is the minimum (total) degree of any real polynomial p such that $\|p - f\|_\infty \leq \varepsilon$, i.e., $|p(x) - f(x)| \leq \varepsilon$ for all $x \in \{-1, 1\}^n$. We use $\widetilde{\text{deg}}(f)$ to denote $\widetilde{\text{deg}}_{1/3}(f)$, and use this to refer to the *approximate degree* of a function without qualification. The choice of $1/3$ is arbitrary, as $\widetilde{\text{deg}}(f)$ is related to $\widetilde{\text{deg}}_\varepsilon(f)$ by a constant factor for any constant $\varepsilon \in (0, 1)$.

Given a Boolean function f , let p be a real polynomial that minimizes $\|p - f\|_\infty$ among all polynomials of degree at most d . Since we work over $x \in \{-1, 1\}^n$, we may assume without loss of generality that p is multilinear with the representation $p(x) = \sum_{|S| \leq d} c_S \chi_S(x)$ where the coefficients c_S are real numbers. Then p is an optimum of the following linear program.

$\begin{array}{ll} \min & \varepsilon \\ \text{such that} & \left f(x) - \sum_{ S \leq d} c_S \chi_S(x) \right \leq \varepsilon \quad \text{for each } x \in \{-1, 1\}^n \\ & c_S \in \mathbb{R} \quad \text{for each } S \leq d \\ & \varepsilon \geq 0 \end{array}$
--

The dual LP is as follows.

$\begin{aligned} \max & \quad \sum_{x \in \{-1, 1\}^n} \phi(x) f(x) \\ \text{such that} & \quad \sum_{x \in \{-1, 1\}^n} \phi(x) = 1 \\ & \quad \sum_{x \in \{-1, 1\}^n} \phi(x) \chi_S(x) = 0 \quad \text{for each } S \leq d \\ & \quad \phi(x) \in \mathbb{R} \quad \text{for each } x \in \{-1, 1\}^n \end{aligned}$
--

Strong LP-duality thus yields the following well-known dual characterization of approximate degree (cf. [50]).

Theorem 14. *Let $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$ be a Boolean function. Then $\widetilde{\deg}_\varepsilon(f) > d$ if and only if there is a polynomial $\phi : \{-1, 1\}^n \rightarrow \mathbb{R}$ such that*

$$\sum_{x \in \{-1, 1\}^n} f(x) \phi(x) > \varepsilon, \quad (1)$$

$$\sum_{x \in \{-1, 1\}^n} |\phi(x)| = 1, \quad (2)$$

and

$$\sum_{x \in \{-1, 1\}^n} \phi(x) \chi_S(x) = 0 \text{ for each } |S| \leq d. \quad (3)$$

If ϕ satisfies Eq. (3), we say ϕ has *pure high degree* d . We refer to any feasible solution ϕ to the dual LP as a *dual polynomial* for f .

3.1.2 One-Sided Approximate Degree

We introduce a relaxed notion of the approximate degree of f which we call the one-sided ε -approximate degree, denoted by $\widetilde{\text{odeg}}_\varepsilon(f)$. This is the least degree of a real polynomial p with that is an ε -one-sided approximation to f , meaning

1. $|p(x) - 1| \leq \varepsilon$ for all $x \in f^{-1}(1)$.
2. $p(x) \leq -1 + \varepsilon$ for all $x \in f^{-1}(-1)$.

That is, we require p to be very accurate on inputs in $f^{-1}(1)$, but only require “one-sided accuracy” on inputs in $f^{-1}(-1)$. We use $\widetilde{\text{odeg}}(f)$ to denote $\widetilde{\text{odeg}}_{1/3}(f)$, and refer to this quantity without qualification as the *one-sided approximate degree* of f .

The primal and dual LPs change in a simple but crucial way if we look at one-sided approximate degree rather than approximate degree. Let $p(x) = \sum_{|S| \leq d} c_S \chi_S(x)$ be a polynomial of degree d for which the ε -one-sided approximate degree of f is attained. Then p is an optimum of the following linear program.

$\begin{aligned} \min & \quad \varepsilon \\ \text{such that} & \quad \left f(x) - \sum_{ S \leq d} c_S \chi_S(x) \right \leq \varepsilon \quad \text{for each } x \in f^{-1}(1) \\ & \quad \sum_{ S \leq d} c_S \chi_S(x) \leq -1 + \varepsilon \quad \text{for each } x \in f^{-1}(-1) \\ & \quad c_S \in \mathbb{R} \quad \text{for each } S \leq d \\ & \quad \varepsilon \geq 0 \end{aligned}$

The dual LP is as follows.

max	$\sum_{x \in \{-1, 1\}^n} \phi(x) f(x)$	
such that	$\sum_{x \in \{-1, 1\}^n} \phi(x) = 1$	
	$\sum_{x \in \{-1, 1\}^n} \phi(x) \chi_S(x) = 0$	for each $ S \leq d$
	$\phi(x) \leq 0$ for each $x \in f^{-1}(-1)$	
	$\phi(x) \in \mathbb{R}$	for each $x \in \{-1, 1\}^n$

We again appeal to strong LP-duality for the following dual characterization of one-sided approximate degree.

Theorem 15. *Let $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$ be a Boolean function. Then $\widetilde{\text{odeg}}_\varepsilon(f) > d$ if and only if there is a polynomial $\phi : \{-1, 1\}^n \rightarrow \mathbb{R}$ such that*

$$\sum_{x \in \{-1, 1\}^n} f(x) \phi(x) > \varepsilon, \quad (4)$$

$$\sum_{x \in \{-1, 1\}^n} |\phi(x)| = 1, \quad (5)$$

$$\sum_{x \in \{-1, 1\}^n} \phi(x) \chi_S(x) = 0 \text{ for each } |S| \leq d, \quad (6)$$

and

$$\phi(x) \leq 0 \text{ for each } x \in f^{-1}(-1). \quad (7)$$

Observe that a feasible solution ϕ to this dual LP is a feasible solution to the dual LP for approximate degree, with the additional constraint that $\phi(x)$ agrees in sign with $f(x)$ whenever $x \in f^{-1}(-1)$. We refer to any such feasible solution ϕ as a dual polynomial for f with *one-sided error*. Dual polynomials with one-sided error have recently played an important role in resolving open problems in communication complexity [19] and resolving the approximate degree of the two-level AND-OR tree [15, 45]. They will play a critical role in our proof of Theorem 1 as well.

Prior work using the dual formulation of one-sided approximate degree exploited the fact that the AND function has one-sided approximate degree equal to its ordinary approximate degree [15, 19, 45]. This fact also plays an important role in the applications of our hardness amplification technique to AND-OR trees (Section 6) and to read-once DNF formulas (Section 7).

Fact 16.

$$\widetilde{\text{odeg}}(\text{AND}_m) = \widetilde{\text{deg}}(\text{AND}_m) = \Omega(\sqrt{m}).$$

Fact 16 can be seen by observing that Nisan and Szegedy's proof that $\widetilde{\text{deg}}(\text{AND}_m) = \Omega(\sqrt{m})$ in fact extends to one-sided approximate degree [39]. Alternatively, it can be directly shown that any dual witness (as defined in Theorem 14) for the fact that $\widetilde{\text{deg}}(\text{AND}_m) = \Omega(\sqrt{m})$ must have one-sided error (cf. [19, Theorem 5.1]).

3.1.3 Approximate Weight

We define the *degree- d ε -approximate weight* of f , $W_\varepsilon(f, d)$, to be the minimum weight of a degree- d polynomial that approximates f pointwise to error ε . Recall that the weight of a polynomial p is the L_1 norm of its coefficients. If $\widetilde{\text{deg}}_\varepsilon(f) > d$, we define $W_\varepsilon(f, d) = \infty$.

For a fixed error parameter ε and degree d , the degree- d ε -approximate weight of a function f is captured by the following optimization problem.

$$\begin{array}{ll}
\min & \sum_{|S| \leq d} |c_S| \\
\text{such that} & \left| f(x) - \sum_{|S| \leq d} c_S \chi_S(x) \right| \leq \varepsilon \quad \text{for each } x \in \{-1, 1\}^n \\
& c_S \in \mathbb{R} \quad \text{for each } |S| \leq d
\end{array}$$

A standard substitution of each term $|c_S|$ in the objective with an auxiliary non-negative variable c'_S , as well as the addition of the constraints $c_S \leq c'_S$ and $-c_S \leq c'_S$ shows that this is in fact a linear program. The dual LP is as follows.

$$\begin{array}{ll}
\max & \sum_{x \in \{-1, 1\}^n} \phi(x) f(x) - \varepsilon \sum_{x \in \{-1, 1\}^n} |\phi(x)| \\
\text{such that} & \left| \sum_{x \in \{-1, 1\}^n} \phi(x) \chi_S(x) \right| \leq 1 \quad \text{for each } |S| \leq d \\
& \phi(x) \in \mathbb{R} \quad \text{for each } x \in \{-1, 1\}^n
\end{array}$$

We thus obtain the following duality theorem.

Theorem 17. *Let $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$ be a Boolean function. Then $W_\varepsilon(f, d) > w$ if and only if there is a polynomial $\phi : \{-1, 1\}^n \rightarrow \mathbb{R}$ such that*

$$\sum_{x \in \{-1, 1\}^n} f(x) \phi(x) - \varepsilon \sum_{x \in \{-1, 1\}^n} |\phi(x)| > w, \quad (8)$$

$$\left| \sum_{x \in \{-1, 1\}^n} \phi(x) \chi_S(x) \right| \leq 1 \text{ for each } |S| \leq d. \quad (9)$$

3.1.4 One-Sided Non-Constant Approximate Weight

To derive our new lower bound on the degree- d threshold weight of read-once DNFs (Corollary 10), we need the following technical variation on approximate weight. Given a polynomial $p(x) = \sum_S c_S \chi_S(x)$, define the *non-constant weight* of p to be the L_1 norm of its coefficients excluding the constant term, i.e., $\sum_{S \neq \emptyset} |c_S|$. We then define the *degree- d one-sided non-constant ε -approximate weight* of f , denoted by $W_\varepsilon^*(f, d)$ to be the minimum non-constant weight of an ε -one-sided approximation to f . Linear programming duality yields the following characterization of $W_\varepsilon^*(f, d)$.

Theorem 18. *Let $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$ be a Boolean function. Then $W_\varepsilon^*(f, d) > w$ if and only if there is a polynomial $\phi : \{-1, 1\}^n \rightarrow \mathbb{R}$ such that*

$$\sum_{x \in \{-1, 1\}^n} f(x) \phi(x) - \varepsilon \sum_{x \in \{-1, 1\}^n} |\phi(x)| > w, \quad (10)$$

$$\left| \sum_{x \in \{-1, 1\}^n} \phi(x) \chi_S(x) \right| \leq 1 \text{ for each } 0 < |S| \leq d, \quad (11)$$

$$\sum_{x \in \{-1, 1\}^n} \phi(x) = 0, \quad (12)$$

$$\phi(x) \leq 0 \text{ for each } x \in f^{-1}(-1). \quad (13)$$

3.1.5 Threshold Degree and Threshold Weight

We say a polynomial $p(x) = \sum_S c_S \chi_S(x)$ with *integer* coefficients is a polynomial threshold function (PTF) for a Boolean function f if p sign-represents f at all Boolean inputs, i.e., if $f(x)p(x) > 0$ for all $x \in \{-1, 1\}^n$. The *threshold degree* of f , $\deg_{\pm}(f)$, is the minimum degree of a PTF for f .

The *threshold weight* $W(f)$ is the minimum weight of any PTF for f . Observe that this definition is only meaningful because the coefficients of any PTF for f are required to be integers, as any positive constant multiple of a PTF for f also sign-represents f . More generally, it is of interest to study the tradeoff between the weight and degree necessary for PTF representations. To this end, we define the *degree- d threshold weight* $W(f, d)$ to be the minimum weight of a degree- d PTF for f . If $\deg_{\pm}(f) > d$, define $W(f, d) = \infty$.

While threshold weight is naturally captured by an *integer* program rather than a linear program, it still admits an important dual characterization, obtained by combining results of Freund [18] and Hajnal et al. [21] (see also [20, 50]).

Theorem 19. *Let $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$ and fix an integer $d \geq \deg_{\pm}(f)$. Then for every probability distribution μ on $\{-1, 1\}^n$,*

$$|\mathbb{E}_{x \sim \mu}[f(x)\chi_S(x)]| \geq \frac{1}{W(f, d)} \text{ for each } |S| \leq d. \quad (14)$$

Moreover, there exists a distribution μ for which

$$|\mathbb{E}_{x \sim \mu}[f(x)\chi_S(x)]| \leq \left(\frac{2n}{W(f, d)}\right)^{1/2} \text{ for each } |S| \leq d. \quad (15)$$

3.2 Relating Degree- d Threshold Weight to High-Error Approximations

In this paper, we will often need to translate lower bounds on $\widetilde{\deg}_{\varepsilon}(f)$ for some function f with ε very close to 1 into lower bounds on the degree- d threshold weight of f . This is possible because degree- d PTFs of weight w are closely related to degree- d pointwise approximations with error $1 - 1/w$. In fact, these notions are essentially equivalent when $w \geq \binom{n}{d}$ [50]. The relationships we will need are formalized in the following lemma.

Lemma 20. *Let $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$ be a Boolean function, and let $w > 0$. Then (1) \Rightarrow (2) \Rightarrow (3).*

(1) $\widetilde{\deg}_{1-\frac{1}{w}}(f) > d$.

(2) $W_{1-\frac{1}{w}}(f, d) > 1$.

(3) $W(f, d) > w$.

Lemma 20 implies that a PTF of degree d and weight w can be transformed into $(1 - 1/w)$ -approximation of degree d . Indeed, the proof will go by way of such a transformation.

Proof. Clearly (1) implies (2), since $W_{1-\frac{1}{w}}(f, d) = \infty$ when $\widetilde{\deg}_{1-\frac{1}{w}}(f) > d$. To show that (2) implies (3), suppose there is a PTF p for f having weight w and degree d . Since p has integer coefficients and is nonzero on Boolean inputs, $|p(x)| \geq 1$ on $\{-1, 1\}^n$. Moreover, $|p(x)| \leq w$ by the weight bound, so the polynomial $\frac{1}{w}p(x)$ is a $(1 - \frac{1}{w})$ -approximation to f with weight 1. \square

Remark: We stress that the converse of Lemma 20 fails badly when $w \ll \binom{n}{d}$. For example, we show in Corollary 10 that for any $d > 0$ there exists a read-once DNF F satisfying $W(F, d) \geq \exp\left(\sqrt{n/d}\right)$. In particular, this yields an exponential lower bound on the degree- d threshold weight of F for any $d = n^{1-\delta}$, with $\delta > 0$ a constant. Yet it follows from a result of Sherstov [48] that $\widetilde{\deg}_{5/3}(F) = O(n^{1/2})$ for any read-once DNF F .

4 Hardness Amplification for Approximate Degree

In this section, we show how to generically transform a circuit f with one-sided approximate degree d into a circuit F with ε -approximate degree d for $\varepsilon = 1 - 2^{-t}$. That is, while f cannot be approximated to error $1/2$ by degree d polynomials, F cannot even be approximated to error $1 - 2^{-t}$ by polynomials of the same degree.

Theorem 1. *Let $f : \{-1, 1\}^m \rightarrow \{-1, 1\}$ be a function with $\widetilde{\text{odeg}}_{1/2}(f) > d$. Let $F : \{-1, 1\}^{mt} \rightarrow \{-1, 1\}$ denote the function $\text{OR}_t(f, \dots, f)$. Then $\widetilde{\text{deg}}_{1-2^{-t}}(F) > d$.*

We remark that it is necessary that the *one-sided* approximated degree of f is large, rather than that just the approximate degree of f is large. Theorem 1 is easily seen to be false with one-sided approximate degree replaced by approximate degree. Consider for example the case where $f = \text{OR}_m$. Then $F = \text{OR}_t(\text{OR}_m, \dots, \text{OR}_m) = \text{OR}_{mt}$. Since $\widetilde{\text{deg}}(\text{OR}_m) = \Omega(\sqrt{m})$, applying Theorem 1 with $\widetilde{\text{deg}}$ in place of odeg would say that $\widetilde{\text{deg}}_{1-2^{-t}}(\text{OR}_{mt}) = \Omega(\sqrt{mt})$. Yet the polynomial $q(y) = \frac{1}{mt}(1/2 - \sum_{i=1}^t \sum_{j=1}^m y_{ij})$ demonstrates that $\widetilde{\text{deg}}_{1-\frac{1}{2mt}}(\text{OR}_{mt}) = 1$ for all values of t . However, Theorem 1 does not apply because the one-sided approximate degree of $f = \text{OR}_m$ is constant.

Proof. Let ψ be a dual polynomial for f with one-sided error whose existence is guaranteed by the assumption that $\widetilde{\text{odeg}}_{1/2}(f) > d$. By Theorem 15, ψ satisfies:

$$\sum_{x \in \{-1, 1\}^m} \psi(x)f(x) > 1/2, \quad (16)$$

$$\sum_{x \in \{-1, 1\}^m} |\psi(x)| = 1, \quad (17)$$

$$\sum_{x \in \{-1, 1\}^m} \psi(x)\chi_S(x) = 0 \text{ for each } |S| \leq d \text{ and} \quad (18)$$

$$\psi(x) \leq 0 \text{ for each } x \in f^{-1}(-1). \quad (19)$$

We will construct a dual solution ζ that witnesses the fact that $\widetilde{\text{deg}}_{1-2^{-t}}(F) > d$. Specifically, ζ must satisfy the three conditions of Theorem 14:

$$\sum_{(x_1, \dots, x_t) \in (\{-1, 1\}^m)^t} \zeta(x_1, \dots, x_t)F(x_1, \dots, x_t) > 1 - 2^{-t}. \quad (20)$$

$$\sum_{(x_1, \dots, x_t) \in (\{-1, 1\}^m)^t} |\zeta(x_1, \dots, x_t)| = 1. \quad (21)$$

$$\sum_{(x_1, \dots, x_t) \in (\{-1, 1\}^m)^t} \zeta(x_1, \dots, x_t)\chi_S(x_1, \dots, x_t) = 0 \text{ for each } |S| \leq d. \quad (22)$$

The construction of ζ is as follows. Let $\mathbf{1}$ denote the all-ones vector. Let $\Psi : \{-1, 1\}^t \rightarrow \{-1, 1\}$ be defined such that $\Psi(\mathbf{1}) = 1/2$, $\Psi(-\mathbf{1}) = -1/2$, and $\Psi(x) = 0$ for all other x . Notice that

$$\sum_{(x_1, \dots, x_t) \in \{-1, 1\}^t} \Psi(x_1, \dots, x_t) = 0 \quad (23)$$

We define $\zeta : (\{-1, 1\}^m)^t \rightarrow \mathbb{R}$ by

$$\zeta(x_1, \dots, x_t) := 2^t \Psi(\dots, \widetilde{\text{sgn}}(\psi(x_i)), \dots) \prod_{i=1}^t |\psi(x_i)|, \quad (24)$$

where $x_i = (x_{i,1}, \dots, x_{i,m})$.

Eq. (24) combines dual functions Ψ and ψ to obtain a dual witness ζ in exactly the same manner as in the works of Sherstov [55, Theorem 3.3] and Lee [34]. The analysis in these works implies without modification that ζ satisfies Equations Eq. (21) and Eq. (22). That is, these works show

Claim 21.

$$\sum_{(x_1, \dots, x_t) \in (\{-1, 1\}^m)^t} |\zeta(x_1, \dots, x_t)| = 1.$$

Claim 22.

$$\sum_{(x_1, \dots, x_t) \in (\{-1, 1\}^m)^t} \zeta(x_1, \dots, x_t) \chi_S(x_1, \dots, x_t) = 0 \text{ for each } |S| \leq d.$$

We provide this analysis in Appendix A for completeness, and here focus on arguing that (20) holds. As we remarked earlier, the properties we exploit to show this are (1) that ψ has one-sided error and (2) that the vector -1 has Hamming distance t from the (unique) input in $\text{OR}_t^{-1}(1)$.

We now prove that (20) holds. Let μ be the distribution on $(\{-1, 1\}^m)^t$ given by $\mu(x_1, \dots, x_t) = \prod_{i=1}^t |\psi(x_i)|$. Since ψ is orthogonal to the constant polynomial, it has expected value 0, and hence the string $(\dots, \widetilde{\text{sgn}}(\psi(x_i)), \dots)$ is distributed uniformly in $\{-1, 1\}^t$ when one samples (x_1, \dots, x_t) according to μ . Observe that

$$\begin{aligned} & \sum_{(x_1, \dots, x_t) \in (\{-1, 1\}^m)^t} \zeta(x_1, \dots, x_t) F(x_1, \dots, x_t) \\ &= 2^t \mathbf{E}_\mu[\Psi(\dots, \widetilde{\text{sgn}}(\psi(x_i)), \dots) \text{OR}_t(\dots, f(x_i), \dots)] \\ &= \sum_{z \in \{-1, 1\}^t} \Psi(z) \left(\sum_{(x_1, \dots, x_t) \in (\{-1, 1\}^m)^t} \text{OR}_t(\dots, f(x_i), \dots) \mu(x_1, \dots, x_t | z) \right), \quad (25) \end{aligned}$$

where $\mu(\mathbf{x} | z)$ denotes the probability of \mathbf{x} under μ , conditioned on $(\dots, \widetilde{\text{sgn}}(\psi(x_i)), \dots) = z$.

Let $A_1 = \{x \in \{-1, 1\}^m : \psi(x) > 0, f(x) = -1\}$ and $A_{-1} = \{x \in \{-1, 1\}^m : \psi(x) < 0, f(x) = 1\}$, so $A_1 \cup A_{-1}$ is the set of all inputs x where the sign of $\psi(x)$ disagrees with $f(x)$. Notice that $\sum_{x \in A_1 \cup A_{-1}} |\psi(x)| < 1/4$ because ψ has correlation $1/2$ with f .

Let λ be the distribution on $\{-1, 1\}^m$ defined by $\lambda(x) = |\psi(x)|$. Then for any bit b ,

$$\Pr_{x \sim \lambda}[f(x) \neq \widetilde{\text{sgn}}(\psi(x)) | \widetilde{\text{sgn}}(\psi(x)) = b] = 2 \sum_{x \in A_b} |\psi(x)|.$$

Therefore, as noted in [55], for any given $z \in \{-1, 1\}^t$, the following two random variables are identically distributed:

- The string $(\dots, f(x_i), \dots)$ when one chooses (\dots, x_i, \dots) from the conditional distribution $\mu(\cdot | z)$.
- The string $(\dots, y_i z_i, \dots)$, where $y \in \{-1, 1\}^t$ is a random string whose i th bit independently takes on value -1 with probability $2 \sum_{x \in A_{z_i}} |\psi(x)| < 1/2$.

Thus, Expression (25) equals

$$\sum_{z \in \{-1, 1\}^t} \Psi(z) \cdot \mathbf{E}[\text{OR}_t(\dots, y_i z_i, \dots)], \quad (26)$$

where $y \in \{-1, 1\}^t$ is a random string whose i th bit independently takes on value -1 with probability 2^{-t} and 1 with probability $1 - 2^{-t}$. We first argue that the term corresponding to $z = \mathbf{1}$ contributes $\Psi(z) = 1/2$ to Expression (26). By Eq. (19), if $f(x) = -1$, then $\psi(x) \leq 0$. This implies that A_1 is empty; that is, if $\widetilde{\text{sgn}}(\psi(x)) = 1$, then it must be the case that $f(x) = 1$. Therefore, for $z = \mathbf{1}$, the y_i 's are all 1 with probability 1, and hence $\mathbf{E}_y[\text{OR}_t(\dots, y_i z_i, \dots)] = \text{OR}_t(\mathbf{1}) = 1$. Thus the term corresponding to $z = \mathbf{1}$ contributes $\Psi(z) \text{OR}_t(z) = 1/2$ to Expression (26) as claimed.

All $z \notin \{\mathbf{1}, -\mathbf{1}\}$ are given zero weight by Ψ and hence contribute nothing to the sum. All that remains is to show that the contribution of the term $z = -\mathbf{1}$ to the sum is $\frac{1}{2}(1 - 2^{-t})$. Since each $y_i = 1$ independently with probability at least $1/2$, and $\text{OR}_t(\dots, -y_i, \dots) = -1$ as long as there is at least one $y_i \neq -1$, we conclude that $\mathbf{E}[\text{OR}_t(\dots, y_i z_i, \dots)] \geq 1 - 2^{-t+1}$. It follows that the term corresponding to $z = -\mathbf{1}$ contributes at least $\frac{1}{2}(1 - 2^{-t+1})$ to the sum. Thus,

$$\sum_{z \in \{-1, 1\}^t} \Psi(z) \cdot \mathbf{E}[\text{OR}_t(\dots, y_i z_i, \dots)] \geq \frac{1}{2} + \frac{1}{2}(1 - 2^{-t+1}) = 1 - 2^{-t}.$$

This completes the proof. \square

Remark: Since the set A_1 within the proof of Theorem 1 is empty, the “combined” dual witness ζ constructed in the proof in fact has one-sided error. Thus, the proof establishes that $\widetilde{\text{odeg}}_{1-2^{-t}}(F) > d$, which is a stronger conclusion than the $\widetilde{\text{deg}}_{1-2^{-t}}(F) > d$ bound appearing in the theorem statement. We chose to state Theorem 1 as an approximate degree lower bound, rather than as a one-sided approximate degree lower bound, for easier comparison with prior work on approximate degree.

5 Lower Bounds for AC^0

In this section, we establish a new lower bound on the one-sided approximate degree of AC^0 . Combining this lower bound with Theorem 1, we establish new lower bounds on accuracy vs. degree tradeoffs for AC^0 . This in turn yields a new upper bound on the discrepancy, and a new lower bound on the threshold weight of AC^0 .

5.1 The One-Sided Approximate Degree of Symmetric Properties

We identify a fairly general criterion under which the one-sided approximate degree of a Boolean function is equal to its approximate degree. This criterion applies to many functions previously studied in the literature, including the AND function, the ELEMENT DISTINCTNESS and COLLISION functions [2,4], and the 2-TO-1 function [9]. Our result applies to Boolean functions corresponding to *symmetric properties*; we refer the reader to Section 2.2.1 for the relevant notation and definitions.

Theorem 2. *Let $f : \{-1, 1\}^m \rightarrow \{-1, 1\}$ be a Boolean function corresponding to a symmetric property ϕ_f of functions $g_x : [N] \rightarrow [R]$. Suppose that for every pair $x, y \in f^{-1}(-1)$, there is a pair of permutations σ on $[R]$ and π on $[N]$ such that $g_x = \sigma \circ g_y \circ \pi$. Then $\widetilde{\text{odeg}}_{\varepsilon}(f) \geq \frac{1}{\log_2 R} \cdot \widetilde{\text{deg}}_{\varepsilon}(f)$ for all $\varepsilon > 0$.*

Proof. Suppose $\widetilde{\text{odeg}}_\varepsilon(f) = d$. Let p be any degree- d one-sided approximation to f with error ε . As described in the proof overview in Section 2.2.1, we show how to transform p into an actual ε -approximation r for f such that $\deg r \leq (\log_2 R) \deg p$. Our transformation from p to r consists of two steps.

In the first step, we turn p into a ‘‘symmetric’’ polynomial $p^{\text{sym}}(x)$ defined below. The following symmetrization lemma shows that the map $p \mapsto p^{\text{sym}}$ increases the degree of p by at most a factor of $\log_2 R$.

Lemma 23. *Let $m = N \cdot \log_2 R$. For $x, y \in \{-1, 1\}^m$, write $y \sim x$ if there is a pair of permutations σ on $[R]$ and π on $[N]$ such that $g_y = \sigma \circ g_x \circ \pi$. Let $p : \{-1, 1\}^m \rightarrow \mathbb{R}$ be a real polynomial. Define*

$$p^{\text{sym}}(x) = \mathbb{E}_{y \sim x}[p(y)].$$

Then $\deg(p^{\text{sym}}) \leq (\log_2 R) \deg(p)$.

The proof of Lemma 23 exploits a result of Ambainis [4] and appears in Appendix B.

We now turn to the second step of our transformation, in which we identify an affine transformation r of p^{sym} that is an actual ε -approximation to f . To this end, we make two further observations about the polynomial p^{sym} .

Claim 24. *If ϕ_f is a symmetric property and p is an ε -one-sided approximation to f , then p^{sym} is also an ε -one-sided approximation to f .*

Claim 25. *Let $S \subseteq \{-1, 1\}^m$. If $x \sim y$ for every pair $x, y \in S$, then p^{sym} is constant on S .*

We first show how these claims together imply the theorem. By Claim 24, p^{sym} is an ε -one-sided approximation to f . By Claim 25, p^{sym} is constant on the set of inputs $f^{-1}(-1)$, where it takes some value $v \leq -1 + \varepsilon$. If $v \geq -1 - \varepsilon$, then p^{sym} is itself an ε -approximation to f and we are done. Otherwise, define the polynomial

$$r(x) = 1 + \frac{2(p^{\text{sym}}(x) - 1)}{|v - 1|}.$$

Then $r(x) = -1$ for all $x \in f^{-1}(-1)$. Moreover, since $|v - 1| \geq 2$, we have $r(x) \in [1 - \varepsilon, 1 + \varepsilon]$ for all $x \in f^{-1}(1)$. Thus r is a ε -approximation to f .

We now proceed to prove Claims 24 and 25.

Proof of Claim 24. Suppose that ϕ_f is a symmetric property and that p is an ε -one-sided approximation to f . Then $p(x) \in [1 - \varepsilon, 1 + \varepsilon]$ for all $x \in f^{-1}(1)$, and $p(x) \leq -1 + \varepsilon$ for all $x \in f^{-1}(-1)$. We focus on some fixed $x \in f^{-1}(1)$; handling inputs in $f^{-1}(-1)$ is entirely analogous. Since ϕ_f is symmetric, we have $y \in f^{-1}(1)$ whenever $y \sim x$. Therefore, $p(y) \in [1 - \varepsilon, 1 + \varepsilon]$ whenever $y \sim x$, so

$$p^{\text{sym}}(x) = \mathbb{E}_{y \sim x}[p(y)] \in [1 - \varepsilon, 1 + \varepsilon].$$

A similar argument holds for $x \in f^{-1}(-1)$, so p^{sym} is an ε -one-sided approximation to f . \square

Proof of Claim 25. Fix a set $S \subseteq \{-1, 1\}^m$, and suppose $x \sim y$ for every $x, y \in S$. Fix some $x^* \in S$, and let $x \in S$ be arbitrary. It suffices to show that $p^{\text{sym}}(x) = p^{\text{sym}}(x^*)$. Write $\sigma \cdot x \cdot \pi$ for the value y for which $g_y = \sigma \circ g_x \circ \pi$. Let σ_x, π_x be a pair of permutations where $x = \sigma_x \cdot x^* \cdot \pi_x$. Note that the map $\sigma \mapsto \sigma \circ \sigma_x$ is a bijection from the symmetric group over $[R]$ to itself; similarly the map $\pi \mapsto \pi_x \circ \pi$ is a bijection from the symmetric group over $[N]$ to itself. Hence, it holds that

$$\begin{aligned} \mathbb{E}_{y \sim x}[p(y)] &= \mathbb{E}_{\sigma, \pi}[p(\sigma \cdot x \cdot \pi)] \\ &= \mathbb{E}_{\sigma, \pi}[p((\sigma \circ \sigma_x) \cdot x^* \cdot (\pi_x \circ \pi))] \\ &= \mathbb{E}_{\sigma, \pi}[p(\sigma \cdot x^* \cdot \pi)] \\ &= \mathbb{E}_{y \sim x^*}[p(y)]. \end{aligned}$$

Thus we have $p^{\text{sym}}(x) = p^{\text{sym}}(x^*)$, showing that p^{sym} is constant on S . \square

The proof of these claims concludes the proof of Theorem 2. \square

5.2 The One-Sided Approximate Degree of AC^0

Prior to our work, the best lower bound on the one-sided approximate degree of an AC^0 function on m variables was $\Omega(\sqrt{m})$ attained by the AND_m function (Fact 16). However, to obtain stronger lower bounds for AC^0 via our hardness amplification technique, we need a constant-depth circuit with polynomially larger one-sided approximate degree. We now exhibit a depth-two circuit having one-sided approximate degree $\tilde{\Omega}(m^{2/3})$. Let N and R be positive integers such that $N \geq R$ and R is a power of 2. We define the ELEMENT DISTINCTNESS function with range R as follows. The function takes $m = N \log_2 R$ bits as input, and interprets its input as N blocks (x_1, \dots, x_N) with each block consisting of $\log_2 R$ bits. Each block is interpreted as a number in the range $[R]$, and the function evaluates to TRUE if and only if all N numbers are distinct.

It is straightforward to check that for $R = \text{poly}(N)$, the ELEMENT DISTINCTNESS function with range R is computed by a CNF formula of polynomial size. Indeed, the function evaluates to TRUE if and only if there is no number $K \in [R]$ for which there is a pair of distinct indices $i, j \in [N]$ such that $x_i = x_j = K$. Thus, the following natural CNF computes ELEMENT DISTINCTNESS (noting that for any fixed K , the inner formula is computed by a bitwise OR):

$$f(x_1, \dots, x_N) = \bigwedge_{K=1}^R \bigvee_{i \neq j} (x_i \neq K) \vee (x_j \neq K).$$

Aaronson and Shi [2] showed that when $R > 3N/2$, the approximate degree of ELEMENT DISTINCTNESS is $\Omega(N^{2/3})$. Ambainis [4] extended the lower bound to the “small-range” case where $R = N$. For the remainder of the paper, we will use the term ELEMENT DISTINCTNESS without qualification to refer to the small-range case.

In the language of properties of functions, the property ϕ_{ED} determines whether a function is one-to-one. That is, $\phi_{\text{ED}}(g) = 1$ if and only if the function $g : [N] \rightarrow [R]$ is injective. Note that ϕ_{ED} is a symmetric property, since injectivity is preserved under permutations of a function’s domain and range. Furthermore, it is straightforward to verify that ELEMENT DISTINCTNESS satisfies the hypothesis of Theorem 2: for any two inputs x, y corresponding to one-to-one functions $g_x, g_y : [N] \rightarrow [R]$, there exist permutations σ, π such that $g_y = \sigma \circ g_x \circ \pi$. Thus, Theorem 2 implies that the one-sided approximate degree of ELEMENT DISTINCTNESS is $\Omega(N^{2/3}/\log R)$.

In a prior version of this work, we gave a different proof of this fact for the small-range case $N = R$ by manipulating a dual witness for the high approximate degree of ELEMENT DISTINCTNESS. We provide this alternative argument in Appendix C. In fact, in the small-range case, the fact the property ϕ_{ED} holds for exactly one input function up to permutation *of the domain only* allows us to prove a stronger one-sided approximate degree lower bound of $\Omega(N^{2/3})$ (i.e., without the loss of a $\log R$ factor that arises from symmetrizing over the range in the proof of Theorem 2).

Corollary 3. *Let $f : \{-1, 1\}^m \rightarrow \{-1, 1\}$ denote the ELEMENT DISTINCTNESS function. Then $\widetilde{\text{odeg}}(f) = \tilde{\Omega}(m^{2/3})$.*

5.3 Accuracy vs. Degree Tradeoffs for AC^0

We are now in a position to prove our new lower bound on “accuracy vs. degree” tradeoffs for pointwise approximating AC^0 functions by polynomials.

Corollary 4. *For every $d > 0$, there is a depth-3 Boolean circuit $F : \{-1, 1\}^n \rightarrow \{-1, 1\}$ of size $\text{poly}(n)$ such that any degree- d polynomial cannot pointwise approximate F to error better than $1 - \exp(-\tilde{\Omega}(nd^{-3/2}))$. In particular, there is a depth-3 circuit F such that any polynomial of degree at most $n^{2/5}$ cannot pointwise approximate F to error better than $1 - \exp(-\tilde{\Omega}(n^{2/5}))$.*

Proof. Let $t = n/d^{3/2}$, and $m = d^{3/2}$. Define $F = \text{OR}_t(f, \dots, f)$ where $f : \{-1, 1\}^m \rightarrow \{-1, 1\}$ computes the ELEMENT DISTINCTNESS problem. The discussion in Section 5.2 implies that f is computed by a depth-2 circuit, and that f has one-sided approximate degree $\tilde{\Omega}(m^{2/3})$. The claim now follows by Theorem 1. \square

5.3.1 On the Tightness of Theorem 1 and Corollary 4

We now argue that the approximate degree lower bound proved in Theorem 1 is essentially tight. In particular, we show that the function F for which Corollary 4 yields a $(1 - \exp(-\tilde{\Omega}(n^{2/5}))$ -error lower bound for approximating polynomials of degree $n^{2/5}$ actually admits a $(1 - \exp(-\tilde{O}(n^{2/5}))$ -approximating polynomial of degree $\tilde{O}(n^{2/5})$.

Our nearly-matching upper bound makes use of a well-known paradigm for constructing low-weight PTFs (and hence, by Lemma 20, low-accuracy pointwise approximations) for composed functions by way of *rational approximations* (see e.g. [12, 49]). Suppose $f : \{-1, 1\}^m \rightarrow \{-1, 1\}$ is pointwise approximated by a rational function in the sense that for every $x \in \{-1, 1\}^m$,

$$\left| f(x) - \frac{p(x)}{q(x)} \right| < \frac{1}{t},$$

where p, q are polynomials of degree d and weight w and $q(x) > 0$ on $\{-1, 1\}^m$. Then observe that the block composition

$$\text{OR}_t(f(x_1), \dots, f(x_t)) = \text{sgn}(1 - t + f(x_1) + \dots + f(x_t)) = \text{sgn}\left(1 - t + \frac{p(x_1)}{q(x_1)} + \dots + \frac{p(x_t)}{q(x_t)}\right).$$

Multiplying $\left(1 - t + \frac{p(x_1)}{q(x_1)} + \dots + \frac{p(x_t)}{q(x_t)}\right)$ by the positive quantity $q(x_1) \cdots q(x_t)$ and clearing denominators yields a PTF for the composed function of degree td and weight at most $w^t(m + tw)$.

We now construct a rational approximation for $f = \text{ELEMENT DISTINCTNESS}$ with the desired properties. Recall from Section 5.2 that ELEMENT DISTINCTNESS on m variables has a CNF representation where the top AND gate has fan-in $s := O(m^3)$ and each OR gate has fan-in $O(\log m)$. It is easy to check that $\text{AND}_s : \{-1, 1\}^s \rightarrow \{-1, 1\}$ admits the rational approximation

$$\frac{ts - 1 + t \sum_{i=1}^s x_i}{ts + 1 + t \sum_{i=1}^s x_i}$$

with error $1/t$, degree $d = 1$, and weight $w = O(st)$. Moreover, each bottom OR gate in the CNF can be computed exactly by a degree $O(\log m)$ polynomial with weight $O(1)$. Composing these constructions yields a rational approximation for ELEMENT DISTINCTNESS with error $1/t$, degree $d = O(\log m) = O(\log t)$ and weight $O(st) = \text{poly}(t)$. Therefore, F has a PTF of degree $\tilde{O}(t)$ and weight $\exp(\tilde{O}(t))$. By the construction of Lemma 20, F also has a $(1 - \exp(-\tilde{O}(t)))$ -approximation of degree $\tilde{O}(t)$. When $t = n^{2/5}$, we obtain a $(1 - \exp(-\tilde{O}(n^{2/5})))$ -approximating polynomial of degree $\tilde{O}(n^{2/5})$ for F as claimed.

5.3.2 A Sharp Threshold in Accuracy-Degree Tradeoffs

The rational approximations developed in the previous section, combined with the lower bound of Theorem 1 and Corollary 4, reveal a “sharp threshold” in the degree required to approximate a particular function F within a given error parameter. Recall that Theorem 1 and Corollary 4 yield a lower bound of $d = \Omega(m^{2/3}/\log m)$ on the ε -approximate degree of $F = \text{OR}_t(f, \dots, f)$, where f is the ELEMENT DISTINCTNESS function on m variables and $\varepsilon = 1 - 2^{-t}$. In the following discussion, consider any $t = d^{1-\Omega(1)}$.

If our goal is to approximate F to within error $(1 - \exp(-\tilde{O}(t)))$, then the rational approximation techniques described in the preceding section yield an approximating polynomial of degree $\tilde{O}(t)$. On the other hand, if we desire even slightly better error of $1 - 2^{-t}$, then our accuracy-degree tradeoff lower bound of Theorem 1 shows that we require degree $d = \omega(t)$. That is, if we demand error that is slightly better than $1 - \exp(-\tilde{O}(t))$, there is an asymptotic jump from $\tilde{O}(t)$ to $\Omega(d)$ in the required degree.

5.4 Discrepancy of AC^0

In this section we prove our new exponentially small upper bound on the discrepancy of a function in AC^0 . Consider a Boolean function $f : X \times Y \rightarrow \{-1, 1\}$, and let $M^{(f)}$ be its communication matrix $M^{(f)} = [f(x, y)]_{x \in X, y \in Y}$. A combinatorial rectangle of $X \times Y$ is a set of the form $A \times B$ with $A \subseteq X$ and $B \subseteq Y$. For a distribution μ over $X \times Y$, the discrepancy of f with respect to μ is defined to be the maximum over all rectangles R of the bias of f on R . That is:

$$\text{disc}_\mu(f) = \max_R \left| \sum_{(x,y) \in R} \mu(x, y) f(x, y) \right|.$$

The discrepancy of f , $\text{disc}(f)$ is defined to be $\min_\mu \text{disc}_\mu(f)$.

Sherstov’s pattern matrix method [50] shows how to generically transform an AC^0 function with high threshold degree or high threshold weight into another AC^0 function with low discrepancy.

Theorem 26 ([50], adapted from Corollary 1.2 and Theorem 7.3). *Let $F : \{-1, 1\}^n \rightarrow \{-1, 1\}$ be given, and define the communication problem $F' : \{-1, 1\}^{4n} \times \{-1, 1\}^{4n} \rightarrow \{-1, 1\}$ by*

$$F'(x, y) = F(\dots, \bigvee_{j=1}^4 (x_{i,j} \wedge y_{i,j}), \dots).$$

Then for every integer $d \geq 0$,

$$\text{disc}(F')^2 \leq \max \left\{ \frac{2n}{W(F, d-1)}, 2^{-d} \right\}.$$

We apply this theorem to the function $F : \{-1, 1\}^n \rightarrow \{-1, 1\}$ of Corollary 4. This function has ε -approximate degree $n^{2/5}$ for $\varepsilon = 1 - 2^{-\tilde{\Omega}(n^{2/5})}$, and hence by Lemma 20 it holds that $W(f, n^{2/5}) = 2^{\tilde{\Omega}(n^{2/5})}$. We thus obtain our new discrepancy upper bound for AC^0 as stated in Corollary 5, restated here for the reader’s convenience.

Corollary 5. *There is a depth-4 Boolean circuit $F' : \{-1, 1\}^n \rightarrow \{-1, 1\}$ with discrepancy $\exp(-\tilde{\Omega}(n^{2/5}))$.*

5.5 Threshold Weight of AC^0

Combing Lemma 20 with Corollary 4 yields Corollary 6, restated here for the reader’s convenience.

Corollary 6. *For every $d > 0$, there is a depth-3 Boolean circuit $F : \{-1, 1\}^n \rightarrow \{-1, 1\}$ of size $\text{poly}(n)$ such that $W(F, d) \geq \exp\left(\tilde{\Omega}(nd^{-3/2})\right)$. In particular, $W(F, n^{2/5}) = \exp\left(\tilde{\Omega}(n^{2/5})\right)$.*

A result of Krause [32] allows us to extend our new degree- d threshold weight lower bound for AC^0 into an $\exp\left(\tilde{\Omega}(n^{2/5})\right)$ degree independent threshold weight lower bound for a related function F' . In Lemma 27 below, we give a slight modification of Krause's original result that is cleaner to apply, and asymptotically recovers Krause's result when the weights under consideration are superpolynomially large. Our restatement admits a new and simple proof based on LP duality that we present in Appendix D.

Lemma 27. *Let $F : \{-1, 1\}^n \rightarrow \{-1, 1\}$ be a Boolean function, and define $F' : \{-1, 1\}^{3n} \rightarrow \{-1, 1\}$ by*

$$F'(x_1, \dots, x_n, y_1, \dots, y_n, z_1, \dots, z_n) := F(\dots, (\bar{z}_i \wedge x_i) \vee (z_i \wedge y_i), \dots).$$

Then for every integer $d \geq 0$,

$$W(F')^2 \geq \min\left\{\frac{W(F, d)}{2n}, 2^d\right\}.$$

Combining Corollary 6 and Lemma 27 yields Corollary 7. This improves over the previous best threshold weight lower bound for AC^0 , which was $\exp\left(\Omega(n^{1/3})\right)$ [33].

Corollary 7. *There is a depth-4 Boolean circuit $F' : \{-1, 1\}^n \rightarrow \{-1, 1\}$ satisfying $W(F') = \exp\left(\tilde{\Omega}(n^{2/5})\right)$.*

Proof. Let F be the circuit of Corollary 4 and let F' be the depth-four circuit obtained by applying Lemma 27 to F . Let $d = n^{2/5}/\log^c n$ for a sufficiently large constant c . Then Corollary 6 implies that $W(F, d) \geq 2n^{2d}$, and hence $W(F') \geq 2^{d/2} = 2^{\tilde{\Omega}(n^{2/5})}$ by Lemma 27. \square

Remark: While the threshold weight bound of Corollary 7 is stated for polynomial threshold functions over $\{-1, 1\}^n$ (i.e., for polynomials that are integer linear combinations of parities), the same threshold weight lower bound also holds for polynomials over $\{0, 1\}^n$, or equivalently, for integer linear combinations of conjunctions. This can be seen as follows.

Given a set $S \subseteq [n]$, let $\text{AND}_S : \{-1, 1\}^n \rightarrow \{-1, 1\}$ denote the AND function restricted to variables in S . Given a sign-representation $p = \sum_S c_S \text{AND}_S$ for F of weight w , let $\sum_S \hat{p}(S) \chi_S$ denote the Fourier representation of p . It is easy to check that the L_1 -norm of the Fourier coefficients of each conjunction AND_S is at most 3, so the weight of the Fourier expansion of p is $w' := \sum_S |\hat{p}(S)| \leq 3w$. However, we cannot simply conclude that $w/3 \geq w' \geq W(f)$ because the coefficients $\hat{p}(S)$ are not necessarily integers.

Nonetheless, note that $|p(x)| \geq 1$ for all $x \in \{-1, 1\}^n$, since p has integer coefficients. That is, p is a sign-representation for f over $\{-1, 1\}^n$ of weight w' and with margin at least 1. It follows by Theorem 19 that $\exp\left(\tilde{\Omega}(n^{2/5})\right) = W(f) \leq 2n(w')^2 = \text{poly}(n, w)$. We conclude that $w = \exp\left(\tilde{\Omega}(n^{2/5})\right)$ as desired.

The same argument shows that all of our lower bounds on degree- d threshold weight proved in this paper hold for PTFs over $\{0, 1\}^n$, in addition to PTFs over $\{-1, 1\}^n$.

6 Lower Bounds for AND-OR Trees

The d -level AND-OR tree (respectively, OR-AND tree) on n variables is a function described by a read-once circuit of depth d consisting of alternating layers of AND gates and OR gates, with the root gate being an AND gate (respectively, an OR gate). We assume throughout this section that all gates have fan-in $n^{1/d}$; for example, the two-level AND-OR tree is a read-once CNF in which all gates have fan-in $n^{1/2}$. The

assumption on the fan-in is not essential to our analysis in this section, which in fact applies to any read-once Boolean circuit such that all gates at any given layer have the same fan-in. We will let $\text{AND-OR}_{d,n}$ (respectively, $\text{OR-AND}_{d,n}$) denote the d -level AND-OR tree (respectively, OR-AND tree) on n variables.

The current authors [15], and independently Sherstov [45], resolved the approximate degree of $\text{AND-OR}_{2,n}$ by proving an optimal $\Omega(n^{1/2})$ lower bound in this case. However, the techniques of [15, 45] break down for the case of depth three or greater; to the best of our knowledge, the best lower bound that follows from prior work is $\Omega(n^{1/4+1/2d})$, which can be derived by combining the depth-two lower bound [15, 45] with an earlier direct-sum theorem of Sherstov [55, Theorem 3.1].

In this section, we extend the methods of our prior work [15] to prove an $\Omega\left(n^{1/2}/\log^{(d-2)/2} n\right)$ lower bound on the approximate degree of $\text{AND-OR}_{d,n}$ for any constant $d > 0$.

Up to a $\log^{(d-2)/2} n$ factor, this matches an upper bound of $O(n^{1/2})$ which was established for AND-OR trees of any depth via a line of work on quantum query algorithms [6, 22, 43]. Specifically, Høyer, Mosca, and De Wolf [22] proved an upper bound of $O(c^{d-1}n^{1/2})$ for some constant c on the approximate degree of any depth- d AND-OR tree in which all gates at any given layer have the same fan-in. Subsequent work by Ambainis et al. [6] established an upper bound of $n^{1/2}2^{O(\sqrt{\log n})}$ for any depth, and further refinements by Reichardt culminated in an $O(n^{1/2})$ upper bound for any depth [43]. A remarkable result of Sherstov [48], on making polynomials robust to noisy inputs, yields a very different proof of Høyer, Mosca, and De Wolf's $O(c^{d-1}n^{1/2})$ upper bound.

Theorem 8. *Let $\text{AND-OR}_{d,n}$ denote the d -level AND-OR tree on n variables. Then $\widetilde{\deg}(\text{AND-OR}_{d,n}) = \Omega\left(n^{1/2}/\log^{(d-2)/2} n\right)$ for any constant $d > 0$.*

Proof. We begin by proving the claimed lower bound for $\text{AND-OR}_{3,n}$ before explaining how to extend the argument to $\text{AND-OR}_{d,n}$ for an arbitrary depth $d > 0$.

Notation. There will be a total of seven intermediate dual witnesses that arise in our construction of a dual witness ψ_7 for $\text{AND-OR}_{3,n}$. We will denote these seven dual witnesses as ψ_1, \dots, ψ_7 . Let $M = n^{1/3}$ denote the fan-in of all gates in $\text{AND-OR}_{3,n}$. Our goal is to construct a dual witness ψ_7 to demonstrate that $\widetilde{\deg}(\text{AND-OR}_{3,n}) = \Omega\left(n^{1/2}/\log^{1/2} n\right)$.

To this end, let ψ_6 be a dual polynomial witnessing the fact that $\widetilde{\text{odeg}}_{.99}(\text{AND}_M) = \Omega(\sqrt{M})$. By Theorem 14, there is some $d_6 = \Omega(\sqrt{M})$ such that ψ_6 satisfies:

$$\sum_{a \in \{-1,1\}^M} \psi_6(a) \text{AND}_M(a) > .99, \quad (27)$$

$$\sum_{a \in \{-1,1\}^M} |\psi_6(a)| = 1, \quad (28)$$

$$\sum_{a \in \{-1,1\}^M} \psi_6(a) \chi_S(a) = 0 \text{ for each } |S| \leq d_6 \text{ and} \quad (29)$$

$$\psi_6(-\mathbf{1}) \leq 0. \quad (30)$$

As stated in the proof outline (Section 2.3), we are ultimately going to construct a function $\psi_5 : \{-1,1\}^{M^2} \rightarrow \mathbb{R}$ that serves as a dual witness to the high approximate degree of $\text{OR-AND}_{2,M^2}$ while having “almost no error on the wrong side”. More formally, we will show

Claim 28. *There exists a dual witness ψ_5 for the fact that the 0.98-approximate degree of $\text{OR-AND}_{2,M^2}$ is at least $\Omega(M/\sqrt{\log n})$ with the following property. If $A_{-1} = \{z \in \{-1, 1\}^{M^2} : \psi_5(z) < 0 \text{ and } \text{OR-AND}_{2,M^2}(z) = 1\}$, then*

$$\sum_{z \in A_{-1}} |\psi_5(z)| \leq n^{-2}.$$

We will then define our final dual witness ψ_7 via

$$\psi_7(x_1, \dots, x_M) := 2^M \psi_6(\dots, \widetilde{\text{sgn}}(\psi_5(x_i)), \dots) \prod_{i=1}^M |\psi_5(x_i)|, \quad (31)$$

where $x_i = (x_{i,1}, \dots, x_{i,M^2})$.

We now prove the existence of ψ_5 (Claim 28) before returning to the analysis of the combined dual witness ψ_7 .

Proof of Claim 28. As discussed in the introduction, the construction of ψ_5 combines our hardness amplification technique (Theorem 1) with the technique of combining dual witnesses in our earlier work [15].

Construction of ψ_5 . Consider the function $\text{OR-AND}_{2,M^2}$. Let $t = 2 \log n$. We view the root OR gate as an OR of ORs, where the top OR has fan-in M/t and the bottom OR gates each have fan-in t . Thus, we are now thinking of the two-level OR-AND tree as a three-level circuit, where the top two levels consist of OR gates, and the bottom level consists of AND gates. Consider the function $F = \text{OR}_t(\text{AND}_M, \dots, \text{AND}_M)$, which allows us to write $\text{OR-AND}_{2,M^2} = \text{OR}_{M/t}(F, \dots, F)$. By our hardness amplification technique, there is a dual witness ψ_3 for the high one-sided approximate degree of F , even with error inverse polynomially close to 1. We will construct ψ_5 by combining ψ_3 with a dual witness ψ_4 for the high approximate degree of $\text{OR}_{M/t}$.

In more detail, applying Theorem 1 to the AND_M function (see Fact 16) yields a dual witness ψ_3 demonstrating that there is some $d_3 = \Omega(\sqrt{M})$ such that $\widetilde{\text{odeg}}_{1-2^{-t}}(F) \geq d_3$ (see the Remark following the proof of Theorem 1). For the case of depth $d = 3$, we may use ψ_3 as a black box. However, to enable induction in the case of general d , we recall that the dual witness ψ_3 was defined via:

$$\psi_3(b_1, \dots, b_t) := 2^t \psi_2(\dots, \widetilde{\text{sgn}}(\psi_1(b_i)), \dots) \prod_{i=1}^M |\psi_1(b_i)|,$$

where $b_i = (b_{i,1}, \dots, b_{i,M})$, ψ_1 was a dual witness to the high one-sided approximate degree of AND_M , and ψ_2 was defined such that $\psi_2(\mathbf{1}) = 1/2$, $\psi_2(-\mathbf{1}) = -1/2$, and ψ_2 evaluates to 0 for all other inputs in $\{-1, 1\}^t$.

By Theorem 15 the dual witness ψ_3 satisfies:

$$\sum_{b \in \{-1, 1\}^{t \cdot M}} \psi_3(b) F(b) > 1 - 2^{-t} = 1 - n^{-2}, \quad (32)$$

$$\sum_{b \in \{-1, 1\}^{t \cdot M}} |\psi_3(b)| = 1, \quad (33)$$

$$\sum_{b \in \{-1, 1\}^{t \cdot M}} \psi_3(b) \chi_S(b) = 0 \text{ for each } |S| \leq d_3 \text{ and} \quad (34)$$

$$\psi_3(b) \leq 0 \text{ for each } b \in F^{-1}(-1). \quad (35)$$

Now let ψ_4 denote a dual witness to the fact that $\widetilde{\text{deg}}_{.99}(\text{OR}_{M/t}) = \Omega(\sqrt{M/t})$. By Fact 16, this dual witness has one-sided error, but on the side opposite from the one we used to define $\widetilde{\text{odeg}}$. Thus there is some $d_4 = \Omega(\sqrt{M/t})$ such that the following equations hold:

$$\sum_{w \in \{-1,1\}^{M/t}} \psi_4(w) \text{OR}_{M/t}(w) > .99, \quad (36)$$

$$\sum_{w \in \{-1,1\}^{M/t}} |\psi_4(w)| = 1, \quad (37)$$

$$\sum_{w \in \{-1,1\}^{M/t}} \psi_4(w) \chi_S(w) = 0 \text{ for each } |S| \leq d_4 \text{ and} \quad (38)$$

$$\psi_4(\mathbf{1}) \geq 0. \quad (39)$$

Finally, we combine the dual witnesses ψ_4 and ψ_3 to obtain the desired function ψ_5 :

$$\psi_5(z_1, \dots, z_{M/t}) := 2^{M/t} \psi_4(\dots, \widetilde{\text{sgn}}(\psi_3(z_i)), \dots) \prod_{i=1}^{M/t} |\psi_3(z_i)|, \quad (40)$$

where $z_i = (z_{i,1}, \dots, z_{i,t \cdot M})$.

Analysis of ψ_5 . The analysis in [15] immediately implies that ψ_5 has L_1 -norm equal to 1, has pure high degree at least $d_3 \cdot d_4 = \Omega(M/\sqrt{t}) = \Omega(M/\sqrt{\log n})$, and that the correlation of ψ_5 with $\text{OR-AND}_{2,M^2}$ is at least $.99 - 2^{-t} \geq .98$. What remains is to show that ψ_5 has “almost no error on the wrong side”. Recall that $A_{-1} = \{z \in \{-1,1\}^{M^2} : \psi_5(z) < 0, \text{OR-AND}_{2,M^2}(z) = 1\}$. We will show that:

$$\sum_{z \in A_{-1}} |\psi_5(z)| \leq n^{-2}. \quad (41)$$

To establish Eq. (41), we first collect some observations. Let $B_{-1} = \{z_i \in \{-1,1\}^{M \cdot t} : \psi_3(z_i) < 0, F(z_i) = 1\}$.

- **Observation 1:** For every $z = (z_1, \dots, z_{M/t}) \in (\{-1,1\}^{t \cdot M})^{M/t}$ in A_{-1} , the following property must hold: $z_i \in B_{-1}$ for every i such that $\psi_3(z_i) < 0$. This holds because $F(z_i) = 1$ for all $i \in \{1, \dots, M/t\}$, since $\text{OR-AND}_{2,M^2}(z) = 1$.
- **Observation 2:** For every $z = (z_1, \dots, z_{M/t}) \in (\{-1,1\}^{t \cdot M})^{M/t} \in A_{-1}$, there must exist a z_i such that $\psi_3(z_i) < 0$. This is because, if $\psi_3(z_i) \geq 0$ for all $i \in \{1, \dots, M/t\}$, then $\psi_5(z)$ agrees in sign with $\psi_4(\mathbf{1}) > 0$ (see Eq. (39)), contradicting the assumption that $z \in A_{-1}$.
- **Observation 3:** Let μ be the distribution on $\{-1,1\}^{M^2}$ defined via: $\mu(z_1, \dots, z_{M/t}) = \prod_{i=1}^{M/t} |\psi_3(z_i)|$. Since ψ_3 is balanced, the string $(\dots, \widetilde{\text{sgn}}(\psi_3(z_i)), \dots)$ is distributed uniformly in $\{-1,1\}^{M/t}$ when one samples $z = (z_1, \dots, z_{M/t})$ according to μ .
- **Observation 4:** Because ψ_3 has correlation $1 - n^{-2}$ with F (see Eq. (32)), the following equation holds:

$$\sum_{z_i \in B_{-1}} |\psi_3(z_i)| \leq \frac{1}{2} n^{-2}.$$

- **Observation 5:** As in the proof of Theorem 1, let $\mu(z|w)$ denote the probability of z under μ , conditioned on $(\dots, \widetilde{\text{sgn}}(\psi_3(z_i)), \dots) = w$. If $z \sim \mu(\cdot|w)$ for some string w where $w_i = -1$, then the probability that $F(z_i) = 1$ when $\widetilde{\text{sgn}}(\psi_3(z_i)) = w_i$ is $2 \sum_{z_i \in B_{-1}} |\psi_3(z_i)|$.

Thus, we may write:

$$\begin{aligned}
\sum_{z \in A_{-1}} |\psi_5(z)| &= \sum_{z \in A_{-1}} 2^{M/t} |\psi_4(\dots, \widetilde{\text{sgn}}(\psi_3(z_i)), \dots)| \prod_i |\psi_3(z_i)| \\
&\leq \sum_{w \in \{-1, 1\}^{M/t}, w \neq \mathbf{1}} |\psi_4(w)| \cdot \Pr_{z \sim \mu(\cdot|w)} [z_i \in B_{-1} \forall i : w_i = -1] \\
&\leq \sum_{w \in \{-1, 1\}^{M/t}} |\psi_4(w)| \cdot n^{-2} \leq n^{-2}.
\end{aligned}$$

Here, the equality holds by definition of ψ_5 (see Eq. (40)), the first inequality holds by Observations 1, 2 and 3, the second inequality holds by Observations 4 and 5, and the fourth inequality holds because the L_1 norm of ψ_4 is 1 (see Eq. (37)). □

Recall that we defined the combined dual witness

$$\psi_7(x_1, \dots, x_M) := 2^M \psi_6(\dots, \widetilde{\text{sgn}}(\psi_5(x_i)), \dots) \prod_{i=1}^M |\psi_5(x_i)|,$$

where ψ_6 is a dual polynomial for the high one-sided approximate degree of the top AND_M function. In the remainder of the proof, we show that ψ_7 is a dual witness for $\text{AND-OR}_{3,n}$.

Bounding the Correlation of ψ_7 with $\text{AND-OR}_{3,n}$. Using Equation Eq. (41), it is possible to adapt the analysis of [15] to show

Claim 29.

$$\sum_x \psi_7(x) \text{AND-OR}_{3,n}(x) > .97.$$

Proof of Claim 29. The idea is to show that

$$\sum_x \psi_7(x) \text{AND-OR}_{3,n}(x) \approx \sum_{a \in \{-1, 1\}^M} \psi_6(a) \text{AND}_M(a) > .99. \quad (42)$$

To this end, let $A_{-1} = \{z \in \{-1, 1\}^{M^2} : \psi_5(z) < 0, \text{OR-AND}_{2,M^2}(z) = 1\}$ as above, and let $A_1 = \{z \in \{-1, 1\}^{2, M^2} : \psi_5(z) \geq 0, \text{OR-AND}_{2,M^2}(z) = -1\}$. Notice that $A_1 \cup A_{-1}$ is the set of all inputs z where the sign of $\psi_5(z)$ disagrees with $\text{OR-AND}_{2,M^2}(z)$. Notice that $\sum_{z \in A_1 \cup A_{-1}} |\psi_5(z)| \leq .01$ because ψ_5 has correlation at least .98 with $\text{OR-AND}_{2,M^2}$.

Let ν be the distribution on $(\{-1, 1\}^{M^2})^M$ given by $\nu(x_1, \dots, x_M) = \prod_{i=1}^M |\nu(x_i)|$. Since ν is orthogonal to the constant polynomial, it has expected value 0, and hence the string $(\dots, \widetilde{\text{sgn}}(\psi_5(x_i)), \dots)$ is distributed uniformly in $\{-1, 1\}^M$ when one samples (x_1, \dots, x_M) according to ν . Let $\nu(x_i|a)$ denote the probability of x_i under ν , conditioned on $(\dots, \widetilde{\text{sgn}}(\psi_5(x_i)), \dots) = a$.

For any given $a \in \{-1, 1\}^M$, the following two random variables are identically distributed:

- The string $(\dots, \text{OR-AND}_{2,M^2}(x_i), \dots)$ when one chooses (\dots, x_i, \dots) from the conditional distribution $\nu(\cdot|a)$.
- The string $(\dots, y_i a_i, \dots)$, where $y \in \{-1, 1\}^M$ is a random string whose i th bit independently takes on value -1 with probability $2 \sum_{x_i \in A_{a_i}} |\nu(x_i)| \leq .02$.

Thus, the left hand side of Expression (42) equals

$$\sum_{a \in \{-1, 1\}^M} \psi_6(a) \cdot \mathbf{E}[\text{AND}_M(\dots, y_i a_i, \dots)], \quad (43)$$

where $y \in \{-1, 1\}^M$ is a random string whose i th bit independently takes on value -1 with probability $2 \sum_{x_i \in A_{a_i}} |\nu(x_i)| \leq .02$.

All $a \neq -\mathbf{1}_M$ can be handled exactly as in [15] and [55] to argue that they contribute at least $(1 - .02)\psi_6(a)$ to the sum. The key property exploited here is that AND_M has low *block-sensitivity* on these points, allowing us to apply the following proposition.

Proposition 30 ([55]). *Let $f : \{-1, 1\}^M \rightarrow \{-1, 1\}$ be a given Boolean function. Let $y \in \{-1, 1\}^M$ be a random string whose i th bit is set to -1 with probability at most $\gamma \in [0, 1]$, and to $+1$ otherwise, independently for each i . Then for every $a \in \{-1, 1\}^M$,*

$$\mathbf{P}_y[f(a_1, \dots, a_M) \neq f(a_1 y_1, \dots, a_M y_M)] \leq 2\gamma \text{bs}_a(f).$$

In particular, since $\text{bs}_a(\text{AND}_M) = 1$ for all $a \neq -\mathbf{1}_M$, Proposition 30 implies that for all $a \neq -\mathbf{1}_M$, and $a = \text{AND}_M$, $\mathbf{P}_y[f(a_1, \dots, a_M) = f(a_1 y_1, \dots, a_M y_M)] \geq 1 - .02$.

We next argue that the term corresponding to $a = -\mathbf{1}_M$ contributes at least $(1 - 2Mn^{-2})\psi_6(a)$ to Expression (43). By Eq. (41) and a union bound, for $a = -\mathbf{1}_M$, the y_i 's are *all* 1 with probability $1 - 2Mn^{-2}$, and hence $\mathbf{E}_y[\text{AND}_M(\dots, y_i a_i, \dots)] \leq (1 - 2Mn^{-2})\text{AND}_M(-\mathbf{1}_M) = -(1 - 2Mn^{-2})$. By Eq. (30), $\widetilde{\text{sgn}}(\psi_6(-\mathbf{1}_M)) = -1$, and thus the term corresponding to $a = -\mathbf{1}_M$ contributes at least $(1 - 2Mn^{-2})\psi_6(a)$ to Expression (26) as claimed. We conclude that $\sum_x \psi_7(x) \text{AND-OR}_{3,n} \geq .97$. \square

Completing the proof for $d = 3$. The proof that ψ_7 has L_1 -norm 1 and has pure high degree at least $d_5 \cdot d_6 = \Omega(n^{1/2}/\log^{1/2}(n))$ is identical to prior work [55] (see also Appendix A). Combined with Claim 29 showing that $\sum_x \psi_7(x) \text{AND-OR}_{3,n} \geq .97$, we conclude that ψ_7 is a dual witness to the fact that $\widetilde{\text{deg}}_{.97}(\text{AND-OR}_{3,n}) = \Omega(n^{1/2}/\log^{1/2}(n))$.

Extending to general d . For ease of exposition, we focus on the case where d is odd; the case of even d is similar. To enable a proof by induction, we will show that

Claim 31. *For d odd, there exists a dual witness ψ_5 showing that the .99-approximate degree of $\text{AND-OR}_{d,n}$ is $\Omega(n^{1/2}/\log^{(d-1)/2}(n))$. Moreover,*

$$\sum_{y \in A_1} |\psi_5(y)| \leq 2n^{-2}, \quad (44)$$

where $A_1 = \{y : \psi_5(y) > 0, \text{AND-OR}_{d,n}(y) = -1\}$.

For d even, the same statement holds for the approximate degree of $\text{OR-AND}_{d,n}$ where we replace Eq. (44) with the corresponding bound on $\sum_{y \in A_{-1}} |\psi_5(y)|$, where $A_{-1} = \{y : \psi_5(y) < 0, \text{OR-AND}_{d,n}(y) = 1\}$.

Eq. (44) intuitively captures the property that ψ'_1 has “almost no error on the wrong side”.

Proof of Claim 31. As a base case of the induction, the dual witness ψ_1 that we used in the case $d = 3$ clearly satisfies the above properties (in fact, ψ_1 had one-sided error, and therefore satisfied an even stronger condition than Eq. (44)).

As suggested by our choice of ψ_5 as the name of the function we want to construct, the inductive case mimics the proof of Claim 28. To emphasize the similarity between this argument and the proof of Claim 28, we will show that assuming the induction hypothesis at level $d - 2$ implies the induction hypothesis at level $d - 1$, for d odd. The case of d even is similar.

To construct a dual witness ψ_5 proving that $\widetilde{\deg}(\text{OR-AND}_{d-1, n^{1-1/d}}) = \Omega(n^{(1-1/d)/2} / \log^{(d-2)/2}(n))$ with “almost no error on the wrong side,” we inductively assume that there exists a dual witness ψ'_1 for the high approximate degree of the function $G = \text{AND-OR}_{d-2, n^{1-2/d}}$ with almost no error on the wrong side. That is, there exists a ψ'_1 and a $d'_1 = \Omega(n^{(1-2/d)/2} / \log^{(d-3)/2}(n))$ such that

$$\sum_{y \in \{-1, 1\}^{n^{1-2/d}}} \psi'_1(y) G(y) > .99, \quad (45)$$

$$\sum_{y \in \{-1, 1\}^{n^{1-2/d}}} |\psi'_1(y)| = 1, \text{ and} \quad (46)$$

$$\sum_{y \in \{-1, 1\}^{n^{1-2/d}}} \psi'_1(y) \chi_S(y) = 0 \text{ for each } |S| \leq d'_1, \quad (47)$$

in addition to having “almost no error on the wrong side.”

Now we set $M = n^{1/d}$, and define $\psi_2, \psi_3, \dots, \psi_5$ exactly as in the case $d = 3$, but with the dual witness ψ'_1 in place of the dual witness ψ_1 . That is, we let $\psi_2 : \{-1, 1\}^t \rightarrow \mathbb{R}$ be defined via $\psi_2(\mathbf{1}) = 1/2$, $\psi_2(-\mathbf{1}) = -1/2$, and $\psi_2(b_i) = 0$ for all other $b_i \in \{-1, 1\}^t$. We define

$$\psi_3(b_1, \dots, b_t) := 2^t \psi_2(\dots, \widetilde{\text{sgn}}(\psi'_1(b_i)), \dots) \prod_{i=1}^M |\psi'_1(b_i)|,$$

where $b_i = (b_{i,1}, \dots, b_{i,M})$. We define ψ_4 to be a dual witness to the fact that $\widetilde{\deg}_{.99}(\text{OR}_{M/t}) = \Omega(\sqrt{M/t})$ for $t = 2 \log n$. We define ψ_5 exactly as in Eq. (40).

The analysis of ψ_5 proceeds as in the proof of Claim 28, with one modification. In the case of $d = 3$, ψ_1 had one-sided error, so we could directly invoke our hardness amplification result (Theorem 1) to conclude that ψ_3 also had one-sided error, as well as correlation $1 - 2^{-t}$ with the target function $\text{OR}_t(G, \dots, G)$. In the case of general d , ψ'_1 does not have one-sided error. However, ψ'_1 “almost” has one-sided error, as formalized by Eq. (44). It is straightforward to modify the proof of Theorem 1 to show though ψ'_1 satisfies a weaker condition than did ψ_1 , the dual witness ψ_3 nonetheless satisfies the following properties.

Let $B_{-1} = \{z_i \in \{-1, 1\}^{n^{1-2/d} \cdot t} : \psi_3(z_i) < 0, \text{OR}_t(G, \dots, G)(z_i) = 1\}$, and let $B_1 = \{z_i \in \{-1, 1\}^{n^{1-2/d} \cdot t} : \psi_3(z_i) > 0, \text{OR}_t(G, \dots, G)(z_i) = -1\}$. Then:

- $\sum_{z_i \in B_{-1}} |\psi_3(z_i)| \leq 2^{-t}$.
- $\sum_{z_i \in B_1} |\psi_3(z_i)| \leq t \cdot 2n^{1-2/d} / n^2$.

That is, ψ_3 has error exponentially small in t on one side, and the error on the other side blows up by at most a factor of t relative to ψ_1 . This permits us to obtain a variant of Eq. (41), namely:

$$\sum_{z \in A_{-1}} |\psi_5(z)| \leq 2tM/n^2, \quad (48)$$

where as above A_{-1} is defined via:

$$A_{-1} = \{z \in \{-1, 1\}^{n^{1-1/d}} : \psi_5(z) < 0, \text{OR-AND}_{d-1, n^{1-1/d}}(z) = 1\}.$$

This completes the induction and the proof. \square

With Claim 31 in hand, we can construct ψ_7 as in the proof of the $d = 3$ case to obtain Theorem 8. That is, we define ψ_6 to be a dual witness to the high one-sided approximate degree of AND_M , and we define ψ_7 exactly as in Eq. (31).

As before, ψ_7 has L_1 -norm 1 and pure high degree at least $d_5 \cdot d_6 = \Omega\left(n^{1/2} / \log^{(d-2)/2}(n)\right)$. Here, $d_5 = \Omega\left(n^{(1-1/d)/2} / \log^{(d-2)/2}(n)\right)$ denotes the pure high degree of ψ_5 and $d_6 = \Omega(M^{1/2})$ denotes the pure high degree of ψ_6 . Finally, the analysis establishing that ψ_7 has high correlation with $\text{AND-OR}_{d,n}$ is the same as in the case of $d = 3$. \square

7 Lower Bounds for Read-Once DNFs

In this section we derive new approximate degree and degree- d threshold weight lower bounds for read-once DNF formulas. The lower bounds we prove are essentially identical to those proved by Beigel [10] and Servedio et al. [44] for the *decision list* ODD-MAX-BIT, which is not computable by a read-once DNF. Our first construction (Corollary 10) yields a degree- d threshold weight lower bound of $2^{\Omega(\sqrt{n/d})}$, matching the lower bound proved by Servedio et al. for the decision list ODD-MAX-BIT. In Section 7.3, we show that this is essentially optimal in the “high-degree” regime where $d = \Omega(n^{1/3})$.

Our second lower bound (Corollary 11) exhibits a DNF with $(1 - 2^{-n/d^2})$ -approximate degree $\Omega(d)$, matching Beigel’s lower bound for ODD-MAX-BIT. As we remarked in Section 2.4, for $d < n^{1/3}$, Corollary 11 is subsumed by Minsky and Papert’s seminal result exhibiting a read-once DNF F with threshold degree $\Omega(n^{1/3})$. However, for $d > n^{1/3}$, it is not subsumed by Minsky and Papert’s result, nor by Corollary 10. While Corollary 10 yields a lower bound on the degree- d threshold weight of read-once DNFs, it does not yield a lower bound on the *approximate-degree* of read-once DNFs. As described in Section 3.2, while $\deg_{1-\frac{1}{w}}(F) > d$ implies that $W(F, d) > w$, the reverse implication does *not* hold when $w \ll \binom{n}{d}$ (and in fact the read-once DNF considered in Corollary 10 is an explicit example of the reverse implication failing badly).

7.1 Extending the Lower Bound of Servedio et al. to Read-Once DNFs

7.1.1 Hardness Amplification for Approximate Weight

We now extend our hardness amplification techniques from approximate degree to approximate weight. This extension forms the technical heart of our proof that the lower bound of Servedio et al. applies to read-once DNFs.

Theorem 9. *Let $f : \{-1, 1\}^m \rightarrow \{-1, 1\}$ be a function with one-sided non-constant approximate weight $W_{3/4}^*(f, d) > w$. Let $F : \{-1, 1\}^{mt} \rightarrow \{-1, 1\}$ denote the function $\text{OR}_t(f, \dots, f)$. Then F has degree- d $(1 - 2^{-t})$ -approximate weight $W_{1-2^{-t}}(F, d) > 2^{-5t}w$.*

Proof. Let ψ be a dual polynomial for f with one-sided error whose existence is guaranteed by the assumption that $W_{3/4}^*(f, d) > w$. Then by Theorem 18, ψ satisfies:

$$\sum_{x \in \{-1,1\}^m} \psi(x)f(x) - \frac{3}{4} \sum_{x \in \{-1,1\}^m} |\psi(x)| > w, \quad (49)$$

$$\left| \sum_{x \in \{-1,1\}^m} \psi(x)\chi_S(x) \right| \leq 1 \text{ for each } 0 < |S| \leq d, \quad (50)$$

$$\sum_{x \in \{-1,1\}^m} \psi(x) = 0, \text{ and} \quad (51)$$

$$\psi(x) \leq 0 \text{ for each } x \in f^{-1}(-1). \quad (52)$$

We will construct a dual solution ζ that witnesses the fact that $W_{1-2^{-t}}(F, d) > 2^{-5t}w$. Specifically, by Theorem 17, ζ must satisfy the following conditions:

$$\sum_{(x_1, \dots, x_t) \in (\{-1,1\}^m)^t} \zeta(x_1, \dots, x_t)F(x_1, \dots, x_t) - (1 - 2^{-t})|\zeta(x_1, \dots, x_t)| > 2^{-5t}w. \quad (53)$$

$$\left| \sum_{(x_1, \dots, x_t) \in (\{-1,1\}^m)^t} \zeta(x_1, \dots, x_t)\chi_S(x_1, \dots, x_t) \right| \leq 1 \text{ for each } |S| \leq d. \quad (54)$$

As before, let $\Psi : \{-1, 1\}^t \rightarrow \{-1, 1\}$ be defined such that $\Psi(\mathbf{1}) = 1/2$, $\Psi(-\mathbf{1}) = -1/2$, and $\Psi(x) = 0$ for all other x , where $\mathbf{1}$ denotes the all-ones vector. We define $\zeta : (\{-1, 1\}^m)^t \rightarrow \mathbb{R}$ by

$$\zeta(x_1, \dots, x_t) := M_t \Psi(\dots, \widetilde{\text{sgn}}(\psi(x_i)), \dots) \prod_{i=1}^t |\psi(x_i)|, \quad (55)$$

where $x_i = (x_{i,1}, \dots, x_{i,m})$ and M_t is a normalization term to be determined later.

We start with Eq. (54) to determine an appropriate choice of M_t . Notice that since Ψ is orthogonal on $\{-1, 1\}^t$ to constant functions, its expected value is 0. Thus, we may write the Fourier representation for Ψ as

$$\Psi(z) = \sum_{\substack{T \subseteq \{1, \dots, t\} \\ T \neq \emptyset}} \hat{\Psi}(T)\chi_T(z)$$

for some real numbers $\hat{\Psi}(T)$. We can thus write

$$\zeta(x_1, \dots, x_t) = M_t \sum_{T \neq \emptyset} \hat{\Psi}(T) \prod_{i \in T} \psi(x_i) \prod_{i \notin T} |\psi(x_i)|.$$

Given a subset $S \subseteq \{1, \dots, t\} \times \{1, \dots, m\}$ with $|S| \leq d$, partition $S = (\{1\} \times S_1) \cup \dots \cup (\{t\} \times S_t)$ where each $S_i \subseteq \{1, \dots, m\}$. Then

$$\begin{aligned} & \sum_{(x_1, \dots, x_t) \in (\{-1,1\}^m)^t} \zeta(x_1, \dots, x_t)\chi_S(x_1, \dots, x_t) \\ &= M_t \sum_{T \neq \emptyset} \hat{\Psi}(T) \prod_{i \in T} \left(\underbrace{\sum_{x_i \in \{-1,1\}^m} \psi(x_i)\chi_{S_i}(x_i)} \right) \prod_{i \notin T} \left(\sum_{x_i \in \{-1,1\}^m} |\psi(x_i)|\chi_{S_i}(x_i) \right). \end{aligned}$$

Since $|S| \leq d$, we have that $|S_i| \leq d$ for every index $i \in \{1, \dots, t\}$. For each set T , each of the underbraced factors is bounded in absolute value by 1 by (50). Writing

$$\|\psi\|_1 := \sum_{x \in \{-1, 1\}^m} |\psi(x)|$$

for notational convenience, we see that

$$\left| \sum_{(x_1, \dots, x_t) \in (\{-1, 1\}^m)^t} \zeta(x_1, \dots, x_t) \chi_S(x_1, \dots, x_t) \right| \leq M_t \sum_{T \neq \emptyset} \hat{\Psi}(T) \|\psi\|_1^{t-|T|} \leq M_t \cdot t 2^{t-1} \|\psi\|_1^{t-1}.$$

Taking $M_t = 2^{-2t} \|\psi\|_1^{1-t}$ gives (54).

We now proceed to verify (53). Let μ be the distribution on $(\{-1, 1\}^m)^t$ given by $\mu(x_1, \dots, x_t) = \|\psi\|_1^{-t} \prod_{i=1}^t |\psi(x_i)|$. Since ψ is orthogonal to the constant polynomial, it has expected value 0, and hence the string $(\dots, \widehat{\text{sgn}}(\psi(x_i)), \dots)$ is distributed uniformly in $\{-1, 1\}^t$ when one samples (x_1, \dots, x_t) according to μ . Observe that

$$\begin{aligned} & \sum_{(x_1, \dots, x_t) \in (\{-1, 1\}^m)^t} \zeta(x_1, \dots, x_t) F(x_1, \dots, x_t) \\ &= M_t \|\psi\|_1^t \mathbf{E}_\mu[\Psi(\dots, \widehat{\text{sgn}}(\psi(x_i)), \dots) \text{OR}_t(\dots, f(x_i), \dots)] \\ &= 2^{-3t} \|\psi\|_1 \sum_{z \in \{-1, 1\}^t} \Psi(z) \left(\sum_{(x_1, \dots, x_t) \in (\{-1, 1\}^m)^t} \text{OR}_t(\dots, f(x_i), \dots) \mu(x_1, \dots, x_t | z) \right), \end{aligned} \quad (56)$$

where $\mu(\mathbf{x}|z)$ denotes the probability of \mathbf{x} under μ , conditioned on $(\dots, \widehat{\text{sgn}}(\psi(x_i)), \dots) = z$.

Let $A_1 = \{x \in \{-1, 1\}^m : \psi(x) > 0, f(x) = -1\}$ and $A_{-1} = \{x \in \{-1, 1\}^m : \psi(x) < 0, f(x) = 1\}$. Then $2 \sum_{x \in A_1 \cup A_{-1}} |\psi(x)| < \frac{1}{4} \|\psi\|_1 - w$ because ψ has correlation at least $w + \frac{3}{4} \|\psi\|_1$ with f .

As before, for any $z \in \{-1, 1\}^t$, the following two random variables are identically distributed:

- The string $(\dots, f(x_i), \dots)$ when one chooses (\dots, x_i, \dots) from the conditional distribution $\mu(\cdot|z)$.
- The string $(\dots, y_i z_i, \dots)$, where $y \in \{-1, 1\}^t$ is a random string whose i th bit independently takes on value -1 with probability $\frac{2}{\|\psi\|_1} \sum_{x \in A_{z_i}} |\psi(x)| < 1/4 - w/\|\psi\|_1$.

Thus, the correlation is

$$2^{-3t} \|\psi\|_1 \sum_{z \in \{-1, 1\}^t} \Psi(z) \cdot \mathbf{E}[\text{OR}_t(\dots, y_i z_i, \dots)], \quad (57)$$

where $y \in \{-1, 1\}^t$ is a random string whose i th bit independently takes on value -1 with probability $2 \sum_{x \in A_{z_i}} |\psi(x)| < 1/4 - w/\|\psi\|_1$. As in the proof of Theorem 1, the one-sided error (52) of the dual witness ψ implies that the input $z = \mathbf{1}$ contributes $\Psi(z) = 1/2$ to Expression (57). All $z \notin \{\mathbf{1}, -\mathbf{1}\}$ are given zero weight by Ψ and hence contribute nothing to the sum. All that remains is to show that the contribution of the term $z = -\mathbf{1}$ to the sum is $\frac{1}{2}(1 - 2^{-2t+1})$. Since each $y_i = 1$ independently with probability at least $3/4 + w/\|\psi\|_1$, and $\text{OR}_t(\dots, -y_i, \dots) = -1$ as long as there is at least one $y_i \neq -1$, we conclude that $\mathbf{E}[\text{OR}_t(\dots, y_i z_i, \dots)] \geq 1 - 2^{-2t+1}$. It follows that the term corresponding to $z = -\mathbf{1}$ contributes at least $\frac{1}{2}(1 - 2^{-2t+1})$ to the sum. Thus,

$$2^{-3t} \|\psi\|_1 \sum_{z \in \{-1, 1\}^t} \Psi(z) \cdot \mathbf{E}[\text{OR}_t(\dots, y_i z_i, \dots)] \geq 2^{-3t} \|\psi\|_1 \left(\frac{1}{2} + \frac{1}{2}(1 - 2^{-2t+1}) \right) = 2^{-3t} (1 - 2^{-2t}) \|\psi\|_1.$$

Since ψ is orthogonal to the constant polynomial by Eq. (51), it has expected value 0, and hence the string $(\dots, \widetilde{\text{sgn}}(\psi(x_i)), \dots)$ is distributed uniformly in $\{-1, 1\}^t$ when one samples (x_1, \dots, x_t) according to μ . Thus,

$$\sum_{(x_1, \dots, x_t) \in (\{-1, 1\}^m)^t} |\zeta(x_1, \dots, x_t)| = 2^{-3t} \|\psi\|_1 \sum_{z \in \{-1, 1\}^t} |\Psi(z)| = 2^{-3t} \|\psi\|_1,$$

Now the left-hand side of Expression (53) is at least

$$2^{-3t}(1 - 2^{-2t})\|\psi\|_1 - (1 - 2^{-t}) \cdot 2^{-3t}\|\psi\|_1 > 2^{-5t}\|\psi\|_1 > 2^{-5t}w,$$

where the last inequality follows from condition (49). This completes the proof. \square

7.1.2 Completing the Proof of Corollary 10

We adapt an argument of Servedio et al. to prove the following one-sided approximate weight lower bound for the function AND_n .

Lemma 32. *Let $d = o(n/\log^2 n)$. Then the function AND_n has one-sided non-constant approximate weight $W_{3/4}^*(\text{AND}_n, d) = 2^{\Omega(n/d)}$.*

Our proof of Lemma 32 follows a symmetrization argument due to Servedio et al. [44]. The key in their proof is the following Markov-type inequality that gives a sharp bound on the derivative of a bounded polynomial in terms of both its degree and weight.

Lemma 33 ([44], Lemma 1). *Let $P : \mathbb{R} \rightarrow \mathbb{R}$ be a degree- d polynomial such that*

1. *The coefficients of P each have absolute value at most w , and*
2. $1/2 \leq \max_{x \in [-1, 1]} |p(x)| \leq R$.

Then $\max_{x \in [-1, 1]} |p'(x)| = O(d \cdot R \cdot \max\{\log W, \log d\})$.

Proof of Lemma 32. Let $p : \mathbb{R}^n \rightarrow \mathbb{R}$ be a real polynomial with degree d and non-constant weight w that has one-sided distance at most $3/4$ from AND_n . Specifically, $p(-\mathbf{1}) \leq -1/4$ and $1/4 \leq p(x) \leq 7/4$ at all other Boolean inputs. We will show that $w = 2^{\Omega(n/d)}$. First observe that if $p(-\mathbf{1}) \leq -7/4$, then the polynomial

$$q(x) = \frac{2(p(x) - 1)}{|p(-\mathbf{1}) - 1|} + 1$$

is a true $(3/4)$ -approximation to AND_n with weight smaller than $w + 1$, so we can assume without loss of generality that p is in fact a $(3/4)$ -approximation to AND_n .

Define the univariate polynomial

$$P(t) := \mathbb{E}_{x \leftarrow \mu_t} [p(x)]$$

where μ_t is the product distribution over $\{-1, 1\}^n$ where each coordinate x_j is independently set to 1 with probability $(1 + t)/2$. Notice that $P(t)$ is obtained from the multivariate expansion of $p(x_1, \dots, x_n)$ by replacing each variable x_i with t . It is readily verified that P satisfies the following properties.

1. $P(-1) = p(-\mathbf{1})$ and $P(1) = p(\mathbf{1})$,
2. $|P(t)| \leq \frac{7}{4}$ for all $t \in [-1, 1]$, and

3. $\deg P \leq \deg p = d$.
4. P has non-constant weight at most w .

By combining properties (1) and (4), we additionally see that the constant term $P(0)$ has absolute value at most $w + \frac{7}{4}$. We can then verify that P satisfies the conditions of Lemma 33.

1. The coefficients of P each have absolute value at most $w + \frac{7}{4}$ and
2. $1/2 \leq \max_{x \in [-1, 1]} |P(t)| \leq \frac{7}{4}$.

Thus we conclude that $|P'(t)| = O(d \max\{\log w, \log d\})$ for $t \in [-1, 1]$. On the other hand, at $t_0 = -1 + 2/n$, we have $\Pr_{x \leftarrow \mu_{t_0}}[x = -1^n] = (1 - \frac{1}{n})^n < 1/e$, so $P(t_0) \geq 1 - \frac{2}{e}$. Since $P(-1) = p(-\mathbf{1}) \leq -\frac{1}{4}$, by the mean value theorem, there is some $t \in [-1, t_0]$ where $P'(t) \geq \frac{n}{4}$. Thus we have $d \max\{\log w, \log d\} = \Omega(n)$, and hence $w = 2^{\Omega(n/d)}$ as long as $d = o(n/\log^2 n)$. \square

Finally, we are in a position to prove Corollary 10, restated here for the reader's convenience.

Corollary 10. *For each $d = o(n/\log^4 n)$, there is a read-once DNF F satisfying $W(F, d) = \exp\left(\Omega\left(\sqrt{n/d}\right)\right)$.*

Proof. Set $m = \alpha\sqrt{nd}$ where α is a constant to be determined later, and let $t = n/m = \Omega(\sqrt{n/d})$. Let $F = \text{OR}_t(\text{AND}_m, \dots, \text{AND}_m)$. By Lemma 32, the inner function AND_m has degree- d one-sided non-constant approximate weight $W_{3/4}^*(\text{AND}_m, d) = 2^{\beta m/d}$ for some constant β . Since $d = o(m/\log^2 m)$, by Theorem 9 the composed function F has degree- d approximate weight

$$W_{1-2^{-t}}(F, d) = 2^{-5t+\beta m/d} = 2^{(-5/\alpha+\beta)\sqrt{n/d}}.$$

Setting $\alpha > 5/\beta$, we get that this approximate weight is greater than 1. By Lemma 20, we have that $W(F, d) > 2^{-t} = 2^{\Omega(\sqrt{n/d})}$. \square

7.2 Extending Beigel's Lower Bound to Read-Once DNFs

Corollary 11. *There is an (explicit) read-once DNF $F : \{-1, 1\}^n \rightarrow \{-1, 1\}$ with $\widetilde{\deg}_{1-2^{-n/d^2}}(F) = \Omega(d)$.*

Proof. Let $m = d^2$, $t = n/d^2$, and $f = \text{AND}_m$. Then Theorem 1 guarantees that

$$\widetilde{\deg}_{1-2^{-t}}(\text{OR}_t(\text{AND}_m, \dots, \text{AND}_m)) > \widetilde{\text{odeg}}(f).$$

By Fact 16, the one-sided approximate degree of f is $\Omega(\sqrt{m})$. This completes the proof. \square

7.3 On the Tightness of Corollaries 10 and 11

In Section 5.3.1, we showed that Corollary 4 is essentially tight by exhibiting a nearly-matching upper bound based on rational approximations. A similar construction shows that any DNF of top fan-in t is computed by a PTF of degree $\tilde{O}(t)$ and weight $\exp\left(\tilde{O}(t)\right)$. This construction immediately shows that Corollary 10 is tight (up to logarithmic factors) for all $d > n^{1/3}$. Indeed, the DNF F for which Corollary 10 demonstrates $W(F, d) \geq \exp(\Omega(\sqrt{n/d}))$ has top fan-in $t = \sqrt{n/d}$, which is less than d for all $d > n^{1/3}$. This construction also reveals a sharp thresholding phenomenon for the read-once DNFs considered in Corollaries 10 and 11 that is similar to the one observed for the depth-three circuit considered in Section 5.3.2.

However, we can provide an alternative construction that demonstrates the tightness of both Corollaries 10 and 11. Specifically, rather than utilizing rational approximation techniques, we can construct a PTF for a read-once DNF by composing a PTF for the top OR gate with low-degree (polynomial, rather than rational) pointwise approximations to each of the individual terms. We provide this construction because of its power to explain why the lower bounds of Corollaries 10 and 11 take their particular forms.

Fix any function $f : \{-1, 1\}^m \rightarrow \{-1, 1\}$, and let $p : \{-1, 1\}^m \rightarrow \{-1, 1\}$ be a polynomial of degree d and weight w such that $|p(x) - f(x)| < 1/t$ for all $x \in \{-1, 1\}^m$. Let $F(x_1, \dots, x_t) = \text{OR}_t(f(x_1), \dots, f(x_t))$. Then for $(x_1, \dots, x_t) \in \{-1, 1\}^{m \cdot t}$, the identity $F(x_1, \dots, x_t) = \text{sgn}(1 - t + \sum_{i=1}^t p(x_i))$ yields a PTF for F of degree at most d and weight at most $tw + t + 1$.

Recall that Corollary 10 yields a lower bound of $W(F, d) = \exp\left(\Omega(\sqrt{n/d})\right)$, where F is the read-once DNF with top fan-in roughly $t = \sqrt{n/d}$ and bottom fan-in roughly $m = \sqrt{nd}$. Servedio et al. [44] showed that for any $d > m^{1/2}$, there is a polynomial p of degree $\tilde{O}(d)$ and weight $\exp\left(\tilde{O}(m/d + \log t)\right) = \exp\left(\tilde{O}\left(\sqrt{n/d}\right)\right)$ that approximates the function AND_m to error $1/t^2$. Hence, as long as $d > n^{1/3}$, the polynomial $1 - t + \sum_{i=1}^t p(x_i)$ is a PTF for F of degree $\tilde{O}(d)$ and weight $\exp\left(\tilde{O}(\sqrt{n/d})\right)$, showing that Corollary 10 is tight up to logarithmic factors.

Similarly, recall that Corollary 11 yields a lower bound of $\widetilde{\text{deg}}_{1-2^{n/d^2}}(F) = \Omega(d)$, where F is the read-once DNF with top fan $t = n/d^2$ and bottom fan-in $m = d^2$. It is well-known that a transformation of the Chebyshev polynomials yields a polynomial p of degree $\tilde{O}(m^{1/2})$ and weight $\exp\left(\tilde{O}(m^{1/2} + \log t)\right)$ that approximates AND_m to error better than $1/t^2$ (see e.g. [30]). Hence, $1 - t + \sum_{i=1}^t p(x_i)$ is a PTF for F of degree $\tilde{O}(d)$ and weight $\exp(\tilde{O}(d + \log t)) = \exp(\tilde{O}(n/d^2))$ when $d < n^{1/3}$. The transformation of Lemma 20 then shows that Corollary 11 is tight up to logarithmic factors in this parameter range.

8 Applications

In this section, we detail applications of the results described above to communication complexity, circuit complexity, and computational learning theory.

8.1 Communication Complexity

Let $f : X \times Y \rightarrow \{-1, 1\}$, where X and Y are finite sets. Consider a two-party communication problem in which Alice is given an input $x \in X$, Bob is given an input $y \in Y$, and their goal is to compute $f(x, y)$ with probability $1/2 + \beta$ for some bias $\beta > 0$. Alice and Bob each have access to an arbitrarily long sequence of private random bits, and the cost $C(P)$ of a protocol P is the worst-case number of bits they must exchange over all inputs $(x, y) \in X \times Y$. Babai et al. [7] defined the *PP communication* model to capture the complexity of computing f with small bias. The PP communication complexity of f , denoted by $\text{PP}(f)$, is the minimum value of $C(P) + \log(1/\beta(P))$ over all protocols P that compute f with positive bias.

It is well known [27] that PP communication is essentially characterized by discrepancy: If $f : \{-1, 1\}^n \times \{-1, 1\}^n \rightarrow \{-1, 1\}$, then $\text{PP}(f) = \Theta(\log(1/\text{disc}(f)) + \log n)$. It follows immediately that our $\exp\left(-\tilde{\Omega}(n^{2/5})\right)$ upper bound on the discrepancy of an AC^0 function f implies an $\tilde{\Omega}(n^{2/5})$ lower bound on $\text{PP}(f)$. The previous best lower bound on $\text{PP}(f)$ for an AC^0 function f was $\Omega(n^{1/3})$ [14, 50].

8.2 Circuit Complexity

Constant-depth circuits of majority gates are known to be surprisingly powerful. Most strikingly, Allender [3] showed that any function in AC^0 can be computed by a depth three circuit of majority gates of quasipolynomial size. This prompted Krause and Pudlák [33] to ask whether every AC^0 function could be computed by depth *two* majority gates of polynomial size. This question was resolved in the negative by Sherstov [51], who exhibited an AC^0 function that cannot be computed even by majority-of-threshold circuits of size $\exp(n^{1/5})$ (later sharpened to $\exp(n^{1/3})$ [50]), and independently by Buhrman, Vereshchagin, and de Wolf [14], who obtained an $\exp(n^{1/3})$ lower bound on the size of majority-of-threshold circuits computing a different AC^0 function.

It is well-known that a discrepancy upper bound for F yields a lower bound on the size of majority-of-threshold circuits computing F [20, 21, 38, 51], and indeed, the circuit lower bounds of [14, 50, 51] are all proved using discrepancy. Through this connection, our discrepancy upper bound of Corollary 5 sharpens the previous lower bounds by yielding a depth-four Boolean circuit F of polynomial size such that any majority-of-threshold circuit computing F requires size $\exp\left(\tilde{\Omega}(n^{2/5})\right)$.

Corollary 34. *There is a depth-four Boolean circuit $F : \{-1, 1\}^n \rightarrow \{-1, 1\}$ of size $\text{poly}(n)$ such that every majority-of-threshold circuit computing F has size $\exp\left(\tilde{\Omega}(n^{2/5})\right)$.*

8.3 Learning Theory

Our results have a number of consequences in computational learning theory. We discuss them below.

Technical Background: The Generalized Winnow Algorithm. The Generalized Winnow algorithm is one of the most powerful known algorithms for online learning [30, 36, 44]. Suppose we are given a concept class \mathcal{C} of functions mapping n -bit inputs to $\{-1, 1\}$, as well as a collection of polynomial-time computable “feature” functions \mathcal{F} . The Generalized Winnow algorithm learns a concept in \mathcal{C} by maintaining as a hypothesis a low-weight linear threshold function of features in \mathcal{F} .

Suppose that each $f \in \mathcal{C}$ has a low-weight linear threshold representation

$$f(x) = \text{sgn} \left(\sum_{h_i \in \mathcal{F}} w_i h_i(x) \right),$$

where each w_i is an integer, and $\sum_i |w_i| \leq W$. A remarkable property of the Generalized Winnow algorithm is that its mistake bound depends only *logarithmically* on the size of the feature set \mathcal{F} , and polynomially on the weight bound W (here the mistake bound refers to the worst-case number of mistakes an online learning algorithm makes over any sequence of examples). Meanwhile, its running time per example is polynomial in the size of the feature set. Standard techniques can be used to transform any online learning algorithm into a PAC learning algorithm whose sample complexity is proportional to the mistake bound.

PAC Learning AC^0 via Generalized Winnow. Valiant famously posed the problem of PAC learning DNF formulas in his original paper [60] introducing the PAC model. The fastest known algorithm for this problem is due to Klivans and Servedio. It is based on linear programming, and takes time $\exp\left(\tilde{O}(n^{1/3})\right)$ [29]. At the core of this algorithm is a fundamental structural result for DNFs: Klivans and Servedio showed that every DNF of size s can be computed by a polynomial threshold function of degree $O(n^{1/3} \log s)$. However, the weight of the PTF arising in this construction can grow doubly-exponentially with n . Klivans and Servedio asked whether it is possible that every polynomial-size DNF has a PTF of degree $\tilde{O}(n^{1/3})$, and weight

$\exp(\tilde{O}(n^{1/3}))$ – an affirmative answer to this question would imply that the Generalized Winnow Algorithm (run over the feature set of all low-degree parities) can also PAC learn DNFs in time $\exp(\tilde{O}(n^{1/3}))$. Such a result would be attractive, as the Generalized Winnow algorithm is substantially simpler than the linear programming algorithm of Klivans and Servedio.

While we do not resolve the question of Klivans and Servedio for DNFs, we do resolve it in the negative for depth-three circuits. In fact, we rule out the possibility of the Generalized Winnow algorithm PAC learning depth-three Boolean circuits in time $\exp(\tilde{O}(n^{2/5}))$ *regardless* of the underlying feature set. That is, our lower bound holds even on feature sets that are not low-degree parities.

Specifically, Corollary 6 implies the following result. The proof is identical to [51, Theorem 8.1] and is omitted for brevity.

Corollary 35. *Let \mathcal{C} denote the concept class of polynomial-size depth-three Boolean circuits. Let $\mathcal{F} = \{h_1, \dots, h_m : \{-1, 1\}^n \rightarrow \{-1, 1\}\}$ be arbitrary Boolean functions such that every $f \in \mathcal{C}$ can be expressed as $f(x) = \text{sgn}(\sum_{i=1}^m w_i h_i(x))$ for some integers w_1, \dots, w_m with $|w_1| + \dots + |w_m| \leq W$. Then $m \cdot W > \exp(\tilde{\Omega}(n^{2/5}))$.*

PAC Learning AC^0 via Boosting. While an $\exp(\tilde{\Omega}(n^{1/3}))$ -time algorithm is known for PAC learning polynomial-size DNF formulas, no $\exp(o(n))$ -time algorithm is known even for learning polynomial-size depth-three Boolean circuits. A natural approach to this problem is as follows. Suppose that every function f in a concept class \mathcal{C} can be computed by a PTF (of arbitrary degree) over $\{0, 1\}^n$ with weight at most W . The well-known *discriminator lemma* of Hajnal et al. [21] implies that under *any* distribution, there is some conjunction (possibly of width $\Omega(n)$) that has correlation at least $1/W$ with f . One can then apply an agnostic learning algorithm for conjunctions (such as the $\exp(\tilde{O}(n^{1/2}))$ -time polynomial regression algorithm of Kalai et al. [24]), combined with standard boosting techniques, to PAC-learn \mathcal{C} in time polynomial in $\max(\exp(\tilde{O}(n^{1/2})), W)$.

Thus, if one could prove an $\exp(\tilde{O}(n^{1/2}))$ upper bound (for PTFs over $\{0, 1\}^n$) on the threshold weight of AC^0 , one would obtain an $\exp(\tilde{O}(n^{1/2}))$ -time algorithm for PAC learning AC^0 . While our $\exp(\tilde{\Omega}(n^{2/5}))$ threshold weight lower bound for AC^0 does not rule out this possibility, it does establish new limitations for this technique. In particular, our threshold weight lower bound implies that even if faster algorithms for agnostically learning conjunctions are discovered, this boosting-based approach to learning AC^0 cannot run in time better than $\exp(\tilde{\Omega}(n^{2/5}))$.

Attribute-Efficient Learning. Attribute-efficient learning is a clean framework that captures the challenging and important problem of learning in the presence of irrelevant information [13]. A class \mathcal{C} of Boolean functions over $\{-1, 1\}^n$ is said to be attribute-efficiently learnable if there is a $\text{poly}(n)$ -time online algorithm that learns any $f \in \mathcal{C}$ with mistake bound polynomial in the representation size of f . For example, the concept class of read-once DNFs that depend on $k \ll n$ of their input variables is attribute-efficiently learnable if there is an online learning algorithm for this class that runs in time $\text{poly}(n)$ per example and achieves mistake bound $\text{poly}(k, \log n)$.

Attribute-efficient learning is a challenging problem, and many simple concept classes are not known to be attribute-efficiently learnable, including decision lists and read-once DNFs. The Generalized Winnow algorithm, run over the feature-space of low-degree parities, marks the best progress toward attribute-efficient learning of these concept classes (see e.g. [30, 44]). Prior to our work, it was unknown whether this approach could learn read-once DNFs depending on k variables in time $\exp(\tilde{O}(n^{1/3}))$ per example and with

mistake bound $\text{poly}(k, \log n)$, as such a guarantee would hold if every read-once DNF on n variables were computed by a polynomial threshold function of degree $\tilde{O}(n^{1/3})$ and weight $\text{poly}(n)$. Corollary 10 rules out this possibility in a very strong sense, as it implies the existence of a read-once DNF that cannot be computed by any PTF of $\text{poly}(n)$ weight, unless the degree is $\tilde{\Omega}(n)$. Similarly, Corollary 4 establishes new limitations on the efficiency of the Generalized Winnow algorithm in the context of attribute-efficient learning of depth-three Boolean circuits.

Acknowledgements. We are grateful to Sasha Sherstov, Robert Špalek, Li-Yang Tan, and the anonymous reviewers for valuable feedback on earlier versions of this manuscript.

References

- [1] S. Aaronson. The polynomial method in quantum and classical computing. In *Proc. of Foundations of Computer Science (FOCS)*, page 3, 2008. Slides available at www.scottaaronson.com/talks/polymeth.ppt.
- [2] S. Aaronson and Y. Shi. Quantum lower bounds for the collision and the element distinctness problems. *Journal of the ACM*, 51(4):595-605, 2004.
- [3] E. Allender. A Note on the Power of Threshold Circuits. *FOCS*, pages 580-584, 1989.
- [4] A. Ambainis. Polynomial degree and lower bounds in quantum complexity: Collision and element distinctness with small range. *Theory of Computing*, 1(1):37-46, 2005.
- [5] A. Ambainis. Polynomial degree vs. quantum query complexity. *J. Comput. Syst. Sci.*, 72(2): pages 220–238, 2006.
- [6] A. Ambainis, A. M. Childs, B. Reichardt, R. Špalek, and S. Zhang. Any AND-OR formula of size N can be evaluated in time $N^{1/2+o(1)}$ on a quantum computer. *SIAM J. Comput.* 39(6): 2513-2530, 2010.
- [7] L. Babai, P. Frankl, and J. Simon. Complexity classes in communication complexity (preliminary version). *FOCS*, pages 337–347, 1986.
- [8] R. Beals, H. Buhrman, R. Cleve, M. Mosca, and R. de Wolf. Quantum lower bound by polynomials. *Journal of the ACM*, 48(4):778-797, 2001.
- [9] P. Beame, W. Machmouchi. The quantum query complexity of AC^0 . *Quantum Information & Computation* 12(7-8): 670-676, 2012.
- [10] R. Beigel. The polynomial method in circuit complexity. *Proc. of the Conference on Structure in Complexity Theory*, pages 82-95, 1993.
- [11] R. Beigel. Perceptrons, PP, and the polynomial hierarchy. *Computational Complexity*, 4:339-349, 1994.
- [12] R. Beigel, N. Reingold, and D. Spielman. PP is closed under intersection. *J. of Comput. and System Sci.*, 50(2):191-202, 1995.
- [13] A. Blum. Learning Boolean functions in an infinite attribute space (extended abstract). *STOC*, pages 64–72. ACM, 1990.
- [14] H. Buhrman, N. K. Vereshchagin, and R. de Wolf. On computation and communication with small bias. *CCC*, pages 24–32, 2007.

- [15] M. Bun and J. Thaler. Dual lower bounds for approximate degree and Markov-Bernstein inequalities. *ICALP*, pages 303-314, 2013.
- [16] A. Chattopadhyay and A. Ada. Multipart communication complexity of disjointness. In *Electronic Colloquium on Computational Complexity (ECCC)*, 08(002), 2008.
- [17] J. Forster, M. Krause, S. V. Lokam, R. Mubarakzjanov, N. Schmitt, and H. Simon. Relations between communication complexity, linear arrangements, and computational complexity. *FSTTCS*, pages 171-182, 2001.
- [18] Y. Freund. Boosting a weak learning algorithm by majority. *Inf. Comput.*, 121(2):256-285, 1995.
- [19] D. Gavinsky and A. A. Sherstov. A separation of NP and coNP in multipart communication complexity. *Theory of Computing*, 6(1):227-245, 2010.
- [20] M. Goldmann, J. Håstad, A. A. Razborov. Majority gates vs. general weighted threshold gates. *Computational Complexity 2*: 277-300, 1992.
- [21] A. Hajnal, W. Maass, P. Pudlák, M. Szegedy, and G. Turán. Threshold circuits of bounded depth. *J. of Comput. and System Sci.*, 46(2):129-154, 1993.
- [22] P. Høyer, M. Mosca, and R. de Wolf. Quantum search on bounded-error inputs. In *Proc. of International Colloquium on Automata, Languages, and Programming*, pages 291–299, 2003.
- [23] A. Kalai, V. Kanade, Y. Mansour. Reliable agnostic learning. *J. Comput. Syst. Sci.* 78(5): 1481-1495, 2012.
- [24] A. Kalai, A. Klivans, Y. Mansour, and R. Servedio. Agnostically learning halfspaces. *SIAM Journal on Computing* 37(6):1777-1805, 2008.
- [25] V. Kanade and J. Thaler. Distribution-independent reliable learning. *CoRR abs/1402.5164*, 2014.
- [26] H. Klauck. On Arthur Merlin games in communication complexity. *CCC*, pages 189–199, 2011.
- [27] H. Klauck. Lower bounds for quantum communication complexity. *FOCS*, pages 288-297, 2001.
- [28] H. Klauck, R. Špalek, and R. de Wolf. Quantum and classical strong direct product theorems and optimal time-space tradeoffs. *SIAM Journal on Computing*, 36(5): 1472-1493, 2007.
- [29] A. R. Klivans and R. A. Servedio. Learning DNF in time $2^{\tilde{O}(n^{1/3})}$. *J. of Comput. and System Sci.*, 68(2):303-318, 2004.
- [30] A. R. Klivans and R. A. Servedio. Toward attribute efficient learning of decision lists and parities. *Journal of Machine Learning Research*, 7:587–602, 2006.
- [31] A. R. Klivans and A. A. Sherstov. Lower bounds for agnostic learning via approximate rank. *Computational Complexity*, 19(4):pages 581–604, 2010.
- [32] M. Krause. On the computational power of Boolean decision lists. *Computational Complexity* 14(4):362-375, 2005.
- [33] M. Krause and P. Pudlák. On the computational power of depth-2 circuits with threshold and modulo gates. *Theor. Comput. Sci.*, 174(1-2): 137-156, 1997.
- [34] T. Lee. A note on the sign degree of formulas. *CoRR abs/0909.4607*, 2009.

- [35] T. Lee and A. Shraibman. Disjointness is hard in the multi-party number-on-the-forehead model. *CCC*, pages 81–91, 2008.
- [36] N. Littlestone. From online to batch learning. *COLT*, pages 269–284, 1989.
- [37] M. L. Minsky and S. A. Papert. *Perceptions: An Introduction to Computational Geometry*. MIT Press, Cambridge, MA., 1969.
- [38] N. Nisan. The Communication Complexity of Threshold Gates. *Combinatorics, Paul Erdos is Eighty*, pages 301-315, 1994.
- [39] N. Nisan and M. Szegedy. On the degree of boolean functions as real polynomials. *Computational Complexity*, 4: pages 301–313, 1994.
- [40] R. O’Donnell and R. Servedio. New degree bounds for polynomial threshold functions. *Combinatorica*, 30(3):327-358, 2010.
- [41] R. Paturi. On the degree of polynomials that approximate symmetric Boolean functions (Preliminary Version). *STOC*, pages 468-474, 1992.
- [42] A. A. Razborov and A. A. Sherstov. The Sign-Rank of AC^0 . *SIAM J. Comput.*, 39(5):1833-1855, 2010.
- [43] B. Reichardt. Reflections for quantum query algorithms. *SODA*, pages 560-569, 2011.
- [44] R. A. Servedio, L.-Y. Tan, and J. Thaler. Attribute-Efficient learning and weight-degree tradeoffs for polynomial threshold functions. *Journal of Machine Learning Research - Proceedings Track*, 23: 14.1-14.19, 2012.
- [45] A. A. Sherstov. Approximating the AND-OR Tree. *Theory of Computing*, 2013.
- [46] A. A. Sherstov Breaking the Minsky-Papert Barrier for constant-depth circuits. *ECCC TR14-009*. To appear in *STOC*, 2014.
- [47] A. A. Sherstov. Communication lower bounds using dual polynomials. *Bulletin of the EATCS*, 95:59-93, 2008.
- [48] A. A. Sherstov. Making polynomials robust to noise. *STOC*, pages 747-758, 2012.
- [49] A. A. Sherstov. Optimal bounds for sign-representing the intersection of two halfspaces by polynomials. *STOC*, pages 523-532, 2010.
- [50] A. A. Sherstov. The pattern matrix method. *SIAM J. Comput.*, 40(6):pages 1969-2000, 2011.
- [51] A. A. Sherstov. Separating AC^0 from depth-2 majority circuits. *SIAM J. Comput.*, 28(6):2113-2129, 2009.
- [52] A. A. Sherstov. Strong direct product theorems for quantum communication and query complexity. *SIAM J. Comput.*, 41(5):1122–1165, 2012.
- [53] A. A. Sherstov. The multiparty communication complexity of set disjointness. *STOC*, pages 525–524, 2012.
- [54] A. A. Sherstov. Communication lower bounds using directional derivatives. *STOC*, pages 921-930, 2013.

- [55] A. A. Sherstov. The intersection of two halfspaces has high threshold degree. *Proc. of Foundations of Computer Science (FOCS)* pages 343-362, 2009. To appear in *SIAM J. Comput. (special issue for FOCS 2009)*
- [56] A. A. Sherstov. The unbounded-error communication complexity of symmetric functions. *Combinatorica*, 31(5):583-614, 2011.
- [57] Y. Shi. Approximating linear restrictions of Boolean functions. Manuscript, 2002. Available online at web.eecs.umich.edu/~shiyiy/mypapers/linear02-j.ps.
- [58] Y. Shi and Y. Zhu. Quantum communication complexity of block-composed functions. *Quantum Information & Computation* 9(5): 444-460, 2009.
- [59] R. Špalek. A dual polynomial for OR. Manuscript, 2008. Available online at: <http://arxiv.org/abs/0803.4516>
- [60] L. Valiant. A theory of the learnable. *Commun. ACM*, 27(11):1134-1142, 1984.

A Final Details of the Proof of Theorem 1

A.1 Proof of Claim 21

Let μ be the distribution on $(\{-1, 1\}^m)^t$ given by $\mu(x_1, \dots, x_t) = \prod_{i=1}^t |\psi(x_i)|$. Since ψ is orthogonal to the constant polynomial, it has expected value 0, and hence the string $(\dots, \widetilde{\text{sgn}}(\psi(x_i)), \dots)$ is distributed uniformly in $\{-1, 1\}^t$ when one samples (x_1, \dots, x_t) according to μ . Thus,

$$\sum_{(x_1, \dots, x_t) \in (\{-1, 1\}^m)^t} |\zeta(x_1, \dots, x_t)| = \sum_{z \in \{-1, 1\}^t} |\Psi(z)| = |\Psi(\mathbf{1})| + |\Psi(-\mathbf{1})| = 1,$$

proving Eq. (21). □

A.2 Proof of Claim 22

We prove that the polynomial ζ defined in Eq. (24) satisfies Eq. (22), reproduced here for convenience.

$$\sum_{(x_1, \dots, x_t) \in (\{-1, 1\}^m)^t} \zeta(x_1, \dots, x_t) \chi_S(x_1, \dots, x_t) = 0 \text{ for each } |S| \leq d. \quad (22)$$

To prove Eq. (22), notice that since Ψ is orthogonal on $\{-1, 1\}^t$ to constant functions, we have the Fourier representation

$$\Psi(z) = \sum_{\substack{T \subseteq \{1, \dots, t\} \\ T \neq \emptyset}} \hat{\Psi}(T) \chi_T(z)$$

for some reals $\hat{\Psi}(T)$. We can thus write

$$\zeta(x_1, \dots, x_t) = 2^t \sum_{T \neq \emptyset} \hat{\Psi}(T) \prod_{i \in T} \psi(x_i) \prod_{i \notin T} |\psi(x_i)|.$$

Given a subset $S \subseteq \{1, \dots, t\} \times \{1, \dots, m\}$ with $|S| \leq d$, partition $S = (\{1\} \times S_1) \cup \dots \cup (\{t\} \times S_t)$ where each $S_i \subseteq \{1, \dots, m\}$. Then

$$\begin{aligned} & \sum_{(x_1, \dots, x_t) \in \{-1, 1\}^m} \zeta(x_1, \dots, x_t) \chi_S(x_1, \dots, x_t) \\ &= 2^t \sum_{T \neq \emptyset} \hat{\Psi}(T) \prod_{i \in T} \underbrace{\left(\sum_{x_i \in \{-1, 1\}^m} \psi(x_i) \chi_{S_i}(x_i) \right)}_{\text{underbraced}} \prod_{i \notin T} \left(\sum_{x_i \in \{-1, 1\}^m} |\psi(x_i)| \chi_{S_i}(x_i) \right). \end{aligned}$$

Since $|S| \leq d$, we have that $|S_i| \leq d$ for every index $i \in \{1, \dots, t\}$. Thus for each set T , at least one of the underbraced factors is zero, as χ_{S_i} is orthogonal to ψ whenever $|S_i| \leq d$.

B Proof of Symmetrization Lemma 23

We now give a proof of Lemma 23, which roughly shows that the symmetrization map $p \mapsto p^{\text{sym}}$ does not increase the degree of p by too much. The notation in the lemma and proof is defined in Section 2.2.1. We also use the shorthand $\sigma \cdot x \cdot \pi$ to denote the boolean vector y for which $g_y = \sigma \circ g_x \circ \pi$.

Lemma 23. *Let $m = N \cdot \log_2 R$. For $x, y \in \{-1, 1\}^m$, write $y \sim x$ if there is a pair of permutations σ on $[R]$ and π on $[N]$ such that $g_y = \sigma \circ g_x \circ \pi$. Let $p : \{-1, 1\}^m \rightarrow \mathbb{R}$ be a real polynomial. Define*

$$p^{\text{sym}}(x) = \mathbb{E}_{y \sim x} [p(y)].$$

Then $\deg(p^{\text{sym}}) \leq (\log_2 R) \deg(p)$.

The proof proceeds in three steps. In the first step, we perform a change of variables showing that $p(x)$ can be written as another polynomial $q(t)$, where $\deg q \leq \deg p$. An input $t \in \{0, 1\}^{N \cdot R}$ to the new polynomial q offers a different representation of a function $g_t : [N] \rightarrow [R]$ as follows: the variable $t_{ij} = 1$ if $g_t(i) = j$, and $t_{ij} = 0$ otherwise.

In the second step, we apply a lemma of Ambainis [4], which shows that q can be partially symmetrized to yield a polynomial Q over yet a different set of variables, again without increasing its degree. This symmetrization yields a polynomial Q whose input now represents a function in a manner invariant under permutations of the function's domain. Specifically, the inputs to the polynomial Q are now variables z_j , where z_j counts the number of $i \in [N]$ for which some function $g_z(i) = j$. Notice that if $g_w = g_z \circ \pi$ for a permutation π , then $w = z$ and hence $Q(w) = Q(z)$.

The third and final step is to symmetrize the polynomial Q once again (without increasing its degree) so that it is invariant under permutations of its input variables z_j . Again interpreting each z_j as the number of i for which some function $g_z(i) = j$, the resulting polynomial Q^{sym} is now invariant under permutations of both the domain *and* codomain of the function g_z .

However, in terms of the original variables x , the new variables z_j are each polynomials of degree $\log_2 R$. Therefore, converting the fully symmetrized polynomial Q^{sym} back into a polynomial p^{sym} over x potentially incurs a $\log_2 R$ factor blow-up in degree.

Proof. Let $p : \{-1, 1\}^m \rightarrow \mathbb{R}$ be a polynomial of degree d . Recall that our proof proceeds in three stages. In the first, we define a new polynomial q over a different set of variables $t \in \{0, 1\}^{N \cdot R}$, and show that $\deg q \leq \deg p$. To this end, define a map $T : \{-1, 1\}^m \rightarrow \{0, 1\}^{N \cdot R}$ by $T_{ij}(x) = 1$ if $g_x(i) = j$, and $T_{ij}(x) = 0$ otherwise. We claim that there is a polynomial $q : \{0, 1\}^{N \cdot R} \rightarrow \mathbb{R}$ of degree d such that

$q(T(x)) = p(x)$ for all $x \in \{-1, 1\}^m$. To see this, write x as a list of blocks $x = (x_1, \dots, x_N)$ where each block has length $\log_2 R$, and let x_{ik} denote the k 'th bit of block x_i . Then

$$x_{ik} = 1 - 2 \sum_{j:j_k=-1} T_{ij}(x),$$

where j_k is the k 'th bit of the binary representation of $j \in [N]$. Hence we can set

$$q(\dots, t_{ij}, \dots) = p \left(\dots, 1 - 2 \sum_{j:j_k=-1} t_{ij}, \dots \right)$$

and have $q(T(x)) = p(x)$, where $\deg q \leq \deg p = d$.

Recall that we can think of the variables t_{ij} themselves as representing functions $g : [N] \rightarrow [R]$ whenever t is the image of a boolean vector under the map T . That is, if $t = T(x)$ for some $x \in \{-1, 1\}^m$, then we can define the function $g_t : [N] \rightarrow [R]$ by $g_t \equiv g_x$. Specifically, $g_t(i)$ is the unique j where $t_{ij} = 1$.

In the second step, we apply a lemma of Ambainis [4] shows that we can symmetrize the polynomial q , again without increasing its degree.

Lemma 36 ([4], Lemma 3.4). *Let $q : \{0, 1\}^{N \cdot R} \rightarrow \mathbb{R}$ be a polynomial of degree d . Then there is a polynomial $Q : \{0, 1, \dots, N\}^R \rightarrow \mathbb{R}$ of degree d such that*

$$Q(\dots, \sum_{i=1}^n t_{ij}, \dots) = \mathbb{E}_\pi [q(t \cdot \pi)]$$

whenever $t = T(x)$ for some $x \in \{-1, 1\}^m$. Here, $t \cdot \pi$ is shorthand for the s for which $g_s = g_t \circ \pi$.

Applying the lemma yields a polynomial Q such that

$$Q(\dots, \sum_{i=1}^n T_{ij}(x), \dots) = \mathbb{E}_\pi [q(T(x) \cdot \pi)] = \mathbb{E}_\pi [p(x \cdot \pi)].$$

For each $j = 1, \dots, R$, let $Z_j(x) = \sum_{i=1}^n T_{ij}(x)$. Notice that $Z_j(x)$ counts the number of inputs $i \in [N]$ for which $g_x(i) = j$. This implies that for any permutation σ on the range $[R]$, we also have that $Z_{\sigma^{-1}(j)}(x)$ counts the number of $i \in N$ for which $(\sigma \circ g_x)(i) = j$. Hence for any fixed σ ,

$$Q(Z_{\sigma^{-1}(1)}(x), \dots, Z_{\sigma^{-1}(R)}(x)) = \mathbb{E}_\pi [p(\sigma \cdot x \cdot \pi)].$$

This observation allows us to complete the third step of the proof, which is symmetrization of the polynomial Q . That is, if we let

$$Q^{\text{sym}}(z) = \mathbb{E}_\sigma [Q(z_{\sigma(1)}, \dots, z_{\sigma(R)})],$$

then

$$Q^{\text{sym}}(Z_1(x), \dots, Z_R(x)) = \mathbb{E}_{\sigma, \pi} [p(\sigma \cdot x \cdot \pi)] = p^{\text{sym}}(x).$$

Notice that since Q is a polynomial of degree d , the symmetrization Q^{sym} clearly also has degree d . To complete the argument, we need to show that each function $Z_j(x)$ is a polynomial of degree $\log_2 R$. To see this, recall that Z_j is a linear combination of the functions T_{ij} , so it suffices to show that T_{ij} has degree $\log_2 R$. The function $T_{ij}(x)$ evaluates to 1 if the block x_i equals to binary representation of j , and evaluates to 0 otherwise. Thus we can write

$$T_{ij}(x) = R \prod_{k=1}^{\log_2 R} (j_k x_{ik} + 1),$$

where x_{ik} is the k 'th bit of block x_i , and j_k is the k 'th bit of the binary representation of j . This expression shows that T_{ij} has degree $\log_2 R$, so the polynomial p^{sym} has degree $d \log_2 R$. \square

C One-Sided Approximate Degree of ELEMENT DISTINCTNESS: Alternative Proof of Corollary 3

Improving on results of Aaronson and Shi [2], Ambainis [4] showed that the ELEMENT DISTINCTNESS problem with small range has approximate degree $\tilde{\Omega}(m^{2/3})$. Recall that the ELEMENT DISTINCTNESS problem on input size $m = N \log_2 N$, where N is a power of 2, takes as input N blocks of length $\log_2 N$ and evaluates to -1 if and only if the blocks are distinct. We will show that there is a dual witness Ψ for the high approximate degree of ELEMENT DISTINCTNESS having one-sided error. Hence, this dual witness actually demonstrates that ELEMENT DISTINCTNESS has high *one-sided* approximate degree.

The idea is that any dual witness for ELEMENT DISTINCTNESS can be “symmetrized” to produce a new dual witness Ψ that is constant on inputs $x \in T$, where T is the set of inputs for which ELEMENT DISTINCTNESS evaluates to true. We then use the fact that Ψ is balanced to argue that the *total* correlation of Ψ with ELEMENT DISTINCTNESS is a constant multiple of the correlation restricted to inputs in T . Since Ψ has positive correlation with ELEMENT DISTINCTNESS, it follows that Ψ must have the correct sign on all inputs in T , as desired.

Formally, let ψ be a dual witness for the fact that $f = \text{ELEMENT DISTINCTNESS}$ has ε -approximate degree $d = \tilde{\Omega}(m^{2/3})$ for some constant ε . By Theorem 14,

$$\sum_{x \in \{-1, 1\}^m} f(x) \psi(x) > \varepsilon, \quad (58)$$

$$\sum_{x \in \{-1, 1\}^m} |\psi(x)| = 1, \quad (59)$$

and

$$\sum_{x \in \{-1, 1\}^m} \psi(x) \chi_S(x) = 0 \text{ for each } |S| \leq d. \quad (60)$$

For any permutation $\sigma \in S^N$, and $x = (x_1, \dots, x_N) \in \{-1, 1\}^m$, define

$$\sigma(x) = (x_{\sigma(1)}, \dots, x_{\sigma(N)}).$$

That is, σ acts on $\{-1, 1\}^m$ by permuting the N blocks of length $\log N$. Observe that for every permutation σ and every $x \in \{-1, 1\}^m$,

$$f(\sigma(x)) = f(x). \quad (61)$$

Now define the symmetrized dual witness

$$\Psi(x) = \mathbb{E}_{\sigma \in S^N} [\psi(\sigma(x))].$$

We will show that Ψ is a dual witness for f with one-sided error by checking the conditions of Theorem 15.

First,

$$\begin{aligned}
\sum_{x \in \{-1,1\}^m} \Psi(x) f(x) &= \mathbb{E}_{\sigma \in S^N} \left[\sum_x \psi(\sigma(x)) f(x) \right] \\
&= \mathbb{E}_{\sigma \in S^N} \left[\sum_x \psi(x) f(x) \right] && \text{by Eq. (61)} \\
&> \epsilon && \text{by (58),}
\end{aligned}$$

verifying (4). Condition (5) is immediate from (59). Condition (6) follows because

$$\sum_{x \in \{-1,1\}^m} \Psi(x) \chi_S(x) = \mathbb{E}_{\sigma \in S^N} \left[\sum_x \psi(x) \chi_{\sigma(S)}(x) \right]$$

where $\sigma(S) = \{\sigma(i) : i \in S\}$ and from (60).

Finally, we check the one-sided error condition (7). We will first show that Ψ is constant on $f^{-1}(-1)$. Let $x^* = (x_1^*, \dots, x_N^*)$ where x_i^* is the binary encoding of i . Since there are only N distinct strings of length $\log N$, $f(x) = -1$ if and only if $x = \sigma_x(x^*)$ for some $\sigma_x \in S^N$. Therefore, if $f(x) = -1$, then

$$\Psi(x) = \mathbb{E}_{\sigma \in S^N} [\psi(\sigma(x))] = \mathbb{E}_{\sigma \in S^N} [\psi((\sigma \circ \sigma_x)(x^*))] = \Psi(x^*),$$

so Ψ is constant on $f^{-1}(-1)$.

By condition (4) it holds that

$$\sum_{x \in f^{-1}(1)} \Psi(x) - \sum_{x \in f^{-1}(-1)} \Psi(x) > \epsilon,$$

and by condition (5) applied to χ_S for $S = \emptyset$ it holds that

$$\sum_{x \in f^{-1}(1)} \Psi(x) + \sum_{x \in f^{-1}(-1)} \Psi(x) = 0.$$

Subtracting the second equation from the first, we conclude that

$$-2 \sum_{x \in f^{-1}(-1)} \Psi(x) > \epsilon.$$

Since Ψ is constant on $f^{-1}(-1)$, this implies that $\Psi(x) < 0$ whenever $x \in f^{-1}(-1)$, proving (7).

D Degree Independent Threshold Weight Bounds via Duality

In this section, we use the dual characterization of threshold weight to give a new proof of a version of Krause's result translating degree- d threshold weight lower bounds for a function F into degree independent threshold weight lower bounds for a related function F' . Specifically, we prove the lemma

Lemma 27. *Let $F : \{-1, 1\}^n \rightarrow \{-1, 1\}$ be a Boolean function, and define $F' : \{-1, 1\}^{3n} \rightarrow \{-1, 1\}$ by*

$$F'(x_1, \dots, x_n, y_1, \dots, y_n, z_1, \dots, z_n) := F(\dots, (\bar{z}_i \wedge x_i) \vee (z_i \wedge y_i), \dots).$$

Then for every integer $d \geq 0$,

$$W(F')^2 \geq \min \left\{ \frac{W(F, d)}{2n}, 2^d \right\}.$$

Proof. By Theorem 19 (condition (14)), it suffices to exhibit a distribution μ' over $\{-1, 1\}^{3n}$ for which

$$|\mathbb{E}_{(x,y,z) \sim \mu'}[F'(x,y,z)\chi_S(x,y,z)]| \leq \max \left\{ \left(\frac{2n}{W(F,d)} \right)^{1/2}, 2^{-d/2} \right\} \text{ for all } S \subseteq \{1, \dots, 3n\}.$$

We construct the distribution μ' as follows. By condition (15) of Theorem 19, there is a probability distribution μ over $\{-1, 1\}^n$ such that

$$|\mathbb{E}_{w \sim \mu}[F(w)\chi_S(w)]| \leq \left(\frac{2n}{W(F,d)} \right)^{1/2} \text{ for each } |S| \leq d. \quad (62)$$

Define $\mu'(x,y,z) = 2^{-2n}\mu(\text{Sel}_z(x,y))$, where $\text{Sel}_z(x,y) = (\dots, (\bar{z}_i \wedge x_i) \vee (z_i \wedge y_i), \dots)$ selects for each index in $[n]$ a bit from either x or y according to z . The distribution μ' has a natural interpretation as follows: it first selects the string z uniformly at random from $\{-1, 1\}^n$. Next, it sets the values of the variables in (x,y) that are selected by z so that they are distributed according to the distribution μ . Finally, it sets the values of the unselected variables in (x,y) uniformly at random.

Note that μ' is indeed a probability distribution, as for every string $w \in \{-1, 1\}^n$, there are exactly 2^{2n} strings (x,y,z) for which $\text{Sel}_z(x,y) = w$. Moreover, this observation allows us to write

$$\mathbb{E}_{(x,y,z) \sim \mu'}[F'(x,y,z)\chi_S(x,y,z)] = 2^{-2n} \sum_{w \in \{-1,1\}^n} F(w)\mu(w) \sum_{(x,y,z): \text{Sel}_z(x,y)=w} \chi_S(x,y,z).$$

Write S as the disjoint union $(\{1\} \times S_1) \cup (\{2\} \times S_2) \cup (\{3\} \times S_3)$ where S_1, S_2, S_3 correspond to indices in x, y, z respectively. Then the expectation becomes

$$2^{-2n} \sum_{z \in \{-1,1\}^n} \chi_{S_3}(z) \underbrace{\sum_{w \in \{-1,1\}^n} F(w)\mu(w) \sum_{(x,y): \text{Sel}_z(x,y)=w} \chi_{S_1}(x)\chi_{S_2}(y)}_{G(z)}$$

Let $G(z)$ denote the underbraced sum.

Suppose there is an index $i \in S_3$ that is not contained in $S_1 \cup S_2$. Then for every $z \in \{-1, 1\}^n$, the string z^i obtained from z by flipping the bit at index i satisfies $\chi_{S_3}(z^i) = -\chi_{S_3}(z)$. On the other hand, for any $(x,y) \in \{-1, 1\}^{2n}$, if we set $x' = (x_1, \dots, x_{i-1}, y_i, x_{i+1}, \dots, x_n)$ and analogously set $y' = (y_1, \dots, y_{i-1}, x_i, y_{i+1}, \dots, y_n)$, then $\text{Sel}_z(x,y) = \text{Sel}_{z^i}(x',y')$. Moreover, because $i \notin S_1 \cup S_2$, it holds that $\chi_{S_1}(x')\chi_{S_2}(y') = \chi_{S_1}(x)\chi_{S_2}(y)$. It follows that $G(z) = G(z^i)$, as each term (x,y) in the underbraced sum defining $G(z)$ is “matched” by term (x',y') in the underbraced sum defining $G(z^i)$. When combined with the fact that $\chi_{S_3}(z^i) = -\chi_{S_3}(z)$, we see that the terms corresponding to z and z^i in the outer sum cancel out, and hence the entire outer sum evaluates to zero. We conclude that for the expectation to be nonzero, we must have $S_3 \subseteq S_1 \cup S_2$, and we assume this holds for the remainder of the proof.

Consider any $i \in S_1$. Then we claim that $G(z) = 0$ whenever z_i selects y_i , i.e., for any z such that $z_i = -1$. This can be seen by another pairing argument: If $\text{Sel}_z(x,y) = w$ but z_i selects y_i , then $\text{Sel}_z(x^i,y) = w$ as well. However, $\chi_{S_1}(x) = -\chi_{S_1}(x^i)$ because $i \in S_1$. This ensures that the innermost sum is zero and hence $G(z) = 0$. The analogous statement holds also for any $i \in S_2$, so for $G(z)$ to be nonzero, it must hold that $z_i = 1$ for all $i \in S_1$ and $z_i = -1$ for all $i \in S_2$. Below, we refer to such a z as a “contributing” z , and all other values of z as “non-contributing”. In particular, we must have $S_1 \cap S_2 = \emptyset$ for z to be contributing.

For any fixed contributing z , it holds that

$$\sum_{(x,y): \text{Sel}_z(x,y)=w} \chi_{S_1}(x)\chi_{S_2}(y) = 2^n \chi_{S_1 \cup S_2}(w).$$

Therefore, it holds that

$$\begin{aligned}
|\mathbb{E}_{(x,y,z)\sim\mu'}[F'(x,y,z)\chi_S(x,y,z)]| &= 2^{-2n} \left| \sum_{z\in\{-1,1\}^n} \chi_{S_3}(z)G(z) \right| \\
&\leq 2^{-n} \sum_{z:G(z)\neq 0} \left| \sum_{w\in\{-1,1\}^n} F(w)\mu(w)\chi_{S_1\cup S_2}(w) \right| \\
&\leq 2^{-|S_1|-|S_2|} \left| \sum_{w\in\{-1,1\}^n} F(w)\mu(w)\chi_{S_1\cup S_2}(w) \right|, \tag{63}
\end{aligned}$$

where inequality (63) used the fact that $G(z) = 0$ for any non-contributing z .

Now we consider two cases for the size of S . First suppose $|S| \leq d$, so in particular, $|S_1 \cup S_2| \leq d$. Then Eq. (62) and inequality (63) implies that

$$|\mathbb{E}_{(x,y,z)\sim\mu'}[F'(x,y,z)\chi_S(x,y,z)]| \leq \left(\frac{2n}{W(f,d)} \right)^{1/2}.$$

Second, suppose that $|S| > d$. We have argued that if $\mathbb{E}_{(x,y,z)\sim\mu'}[F'(x,y,z)\chi_S(x,y,z)] \neq 0$, then $S_3 \subseteq S_1 \cup S_2$. Hence, it must be the case that $|S_1| + |S_2| \geq |S|/2 > d/2$. Therefore, inequality (63) implies that $\mathbb{E}_{(x,y,z)\sim\mu'}[F'(x,y,z)\chi_S(x,y,z)] \leq 2^{-d/2}$. This completes the proof. □